

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C12N 15/12, 5/10, C12Q 1/68, A01K 67/00, C12N 15/00, C07K 16/28, 14/705</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/08157</b> <b>(43) International Publication Date:</b> 17 February 2000 (17.02.00)
<b>(21) International Application Number:</b> PCT/US99/17823 <b>(22) International Filing Date:</b> 6 August 1999 (06.08.99) <b>(30) Priority Data:</b> 60/095,835 7 August 1998 (07.08.98) US <b>(71) Applicant (for all designated States except US):</b> AXYS PHARMACEUTICALS, INC. [US/US]; 180 Kimball Way, South San Francisco, CA 94080 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> LAUBERT, Boris [US/US]; 4550 Bancroft Street #4, San Diego, CA 92116 (US). CARDOSO, Gizela [US/US]; 721 North Main Street, Brockton, Maine 02401 (US). HU, Ping [US/US]; 5807 Folkstone Road, Bethesda, MD 20817 (US). MILLER, Andrew, P. [US/US]; 3271 Countryside Drive, San Mateo, CA 94403 (US). BUCKLER, Alan, J. [US/US]; 2315 Lagoon View Drive, Cardiff, CA 92007 (US). <b>(74) Agent:</b> SHERWOOD, Pamela, J.; Bozicevic, Field & Francis LLP, Suite 200, 285 Hamilton Avenue, Palo Alto, CA 94301 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> HUMAN ANION TRANSPORTER GENES ATNOV  <b>(57) Abstract</b>  Methods for isolating ATnov genes are provided. The ATnov nucleic acid compositions find use in identifying homologous or related proteins and the DNA sequences encoding such proteins; in producing compositions that modulate the expression or function of the protein; and in studying associated physiological pathways. In addition, modulation of the gene activity in vivo is used for prophylactic and therapeutic purposes, such as identification of cell type based on expression, and the like.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## HUMAN ANION TRANSPORTER GENES

## INTRODUCTION

## Background

5           Endo- and xenobiotics are typically cleared from mammals via the liver, the primary site of drug metabolizing enzymes. Charged compounds, either endogenously or exogenously derived, are taken up by hepatocytes across the basolateral membrane, appropriately metabolized by the liver enzymes, trafficked through the cell, and then excreted across the canalicular membrane into the bile. These four steps are important in  
10       determining a patient's response to pharmaceutical agents.

          Generation of bile flow is a regulated, ATP-dependent process and depends on the coordinated action of a number of transporter proteins in the sinusoidal and canalicular domains of the hepatocyte. Dysfunction of any of these proteins leads to retention of substrates, with conjugated hyperbilirubinemia or cholestasis as a result. In recent years  
15       many of the transport proteins involved in bile formation have been identified, cloned, and functionally characterized. The hepatocyte sinusoidal membrane contains transport proteins for the hepatic uptake of organic anions and cations and for the uptake of bile acids.

          The Na<sup>+</sup>-independent organic anion transporter, OATP, resides on the basolateral surface of hepatocytes and mediates the uptake of a large number of amphipathic  
20       substrates, such as bromosulfophalein, bile acids, estrogen conjugates, neutral steroids, organic cations, cardiac glycosides, and peptidomimetic drugs. The human organic anion transporter, OATP, is expressed in multiple tissues, including brain, lung, liver, kidney, and testes, while a rat homolog of OATP is expressed only in liver and kidney (Bergwerk et al. (1996) Am. J. Physiol. 271: G231-G238). A prostaglandin transporter, hPGT, which shares  
25       significant homology with these organic anion transporters, is abundantly expressed (Lu et al. (1996) J. Clin. Invest. 98: 1142-1149).

          Variations in transporter sequences may alter the kinetic properties of the protein. For example, inefficient clearance of substrates would result in an increased biological half-life, where drugs have an increased half-life and drug levels approach or reach toxic  
30       thresholds. Alternatively, over-efficient clearance of substrates could reduce the biological effectiveness of a drug. The identification of novel genes within these pathways provides additional targets for pharmacogenetic analysis, as well as a more thorough understanding of the biological process of drug clearance.

### Relevant Literature

The molecular and functional characterization of an organic anion transporting polypeptide cloned from human liver, OATP, is described by Kullak-Ublick et al. (1995) Gastroenterology 109:1274-1282. Other cloned transporter genes are described by Noe et al. (1997) Proc. Natl. Acad. Sci. 94:10346-10350; and Jacquemin et al. (1994) Proc. Natl. Acad. Sci. 91:133-137.

The role of organic cation transporters in intestine, kidney, liver, and brain is reviewed by Koepsell (1998) Annu Rev Physiol 60:243-266. Canalicular multispecific organic anion transporter and the disposal of endo- and xenobiotics is reviewed by Elferink and Jansen (1994) Pharmac. Ther. 64:77-97.

Public EST sequences having sequence similarity with ATnov nucleic acids include: Genbank accessions nos. N49902 (ATnov2); N50005 (ATnov2); H62927 (ATnov3); H62893 (ATnov3); R29414 (ATnov3); AA382692 (ATnov3); T73863 (ATnov3); T74263 (ATnov3); T55488 (ATnov3).

### SUMMARY OF THE INVENTION

Isolated nucleotide compositions and sequences are provided for ATnov genes. The ATnov nucleic acid compositions find use in identifying homologous or related genes; in producing compositions that modulate the expression or function of its encoded proteins; for gene therapy; mapping functional regions of the proteins; and in studying associated physiological pathways. In addition, modulation of the gene activity in vivo is used for prophylactic and therapeutic purposes, such as treatment of anion transporter defects, identification of cell type based on expression, and the like.

### DESCRIPTION OF THE SPECIFIC EMBODIMENTS

Nucleic acid compositions encoding ATnov anion transporters are provided. They are used in identifying homologous or related genes; in producing compositions that modulate the expression or function of the encoded proteins; for gene therapy; mapping functional regions of the proteins; and in studying associated physiological pathways. The ATnov gene products are members of the anion transporter gene family, and have high degrees of homology at the amino acid level with known anion transporters.

## CHARACTERIZATION OF ATNOV

The sequence data predict that the provided ATnov genes encode anion transporters. Characterization of organic ion transport across the cell membrane, in terms of substrates, binding and transport kinetics, is an important aspect of ATnov biology. A  
5 substrate, as used herein, is a chemical entity that is transported by an ATnov polypeptide, usually under normal physiological conditions. Substrates can be either endogenous substrates, i.e. substrates normally found within the natural environment, such as bile salts, or exogenous, i.e. substrates that are not normally found within the natural environment.

Substrate screening assays are used to determine the kinetics of a ATnov protein or  
10 peptide fragment on a substrate. Many suitable assays are known in the art, including the use of primary or cultured cells, genetically modified cells (e.g., where DNA encoding the ATnov polymorphism to be studied is introduced into the cell within an artificial construct), cell-free systems, e.g. recombinantly produced enzymes in a suitable buffer, or in animals, including human clinical trials (see, e.g. (1995) Burchell et al. Life Sci. 57:1819-1831,  
15 specifically incorporated herein by reference). Where genetically modified cells are used, since most cell lines do not express ATnov activity (liver cells lines being the exception), introduction of artificial construct for expression of the ATnov polymorphism into many human and non-human cell lines does not require additional modification of the host to inactivate endogenous ATnov expression/activity. Clinical trials may monitor serum, urine,  
20 etc. levels of the substrate or its metabolite(s).

Full length ion transporter cDNAs may be combined with proper vectors to form expression constructs of each individual transporter. Functional analyses of expressed transporters can be performed in heterologous systems, or by expression in mammalian cell lines. For expression analyses in heterologous systems such as *Xenopus* oocytes, synthetic  
25 mRNA is made through in vitro transcription of each transporter construct. mRNA is then injected into prepared oocytes and the cells allowed to express the transporter for several days. Candidate substrates may be labeled to provide a means of following movement across the membrane. Similarly, the requirements of a transporter for ATP, Na<sup>+</sup>, etc. may be assessed. For an example of these techniques, see Kullak-Ublick et al. (1997)  
30 Gastroenterology 113(4):1295-1305.

Heterologous or mammalian cell lines expressing the novel transporters can be used to characterize small molecules and drugs that interact with the transporter. The same

experiments can be used to assay for novel compounds that interact with the expressed transporters.

#### ATNOV NUCLEIC ACID COMPOSITIONS

5           As used herein, the term "ATnov" is generically used to refer to any one of the provided nucleotide sequences as set forth in the SEQLIST. Of particular interest are the sequences, including polymorphisms, of ATnov3.1 and ATnov3.2. These sequences are provided as SEQ ID NO:3 (ATnov3.1), SEQ ID NO:5 (ATnov3.1), SEQ ID NO:7 (ATnov3.2) and SEQ ID NO:9 (ATnov3.2). The encoded polypeptides are provided as SEQ ID NO:4,  
10   6, 8 and 10, respectively. The polymorphic variants are set forth in the sequences listings. These include a G or A polymorphism at nucleotide 487, resulting in an amino acid change of asp to asn. There is a polymorphism of C or T at nucleotide 670, which is silent with respect to the encoded polypeptide. A frameshift variant is found in the poly T stretch between positions 1705 and 1710, where the sequence contains either 5T or 6T. The 5T  
15   polymorphism results in a truncated polypeptide product of 542 amino acids (SEQ ID NO:4 and SEQ ID NO:8), while the 6T polymorphism encodes the full-length protein of 591 amino acids.

Also of interest are the genetic sequences of SEQ ID NO:1 (ATnov1) and SEQ ID NO:2 (ATnov2).

20           Where a specific ATnov sequence is intended, the numerical designation will be added. Nucleic acids encoding ATnov anion transporters may be cDNA or genomic DNA or a fragment thereof. The term "ATnov gene" shall be intended to mean the open reading frame encoding any of the provided ATnov polypeptides, introns, as well as adjacent 5' and 3' non-coding nucleotide sequences involved in the regulation of expression, up to about 20  
25   kb beyond the coding region, but possibly further in either direction. The gene may be introduced into an appropriate vector for extrachromosomal maintenance or for integration into a host genome.

Novel nucleic acid compositions of the invention of particular interest comprise a sequence set forth in SEQ ID NO:1, 2, 3, 5, 7, 9 or an identifying sequence thereof. An  
30   "identifying sequence" is a contiguous sequence of residues at least about 10 nt to about 20 nt in length, usually at least about 50 nt to about 100 nt in length, that uniquely identifies a nucleic acid sequence, e.g., exhibits less than 90%, usually less than about 80% to about 85% sequence identity to any contiguous nucleotide sequence of more than about 20 nt.

Thus, the subject novel nucleic acid compositions include full length cDNAs or mRNAs that encompass an identifying sequence of contiguous nucleotides from SEQ ID NO:1, 2, 3, 5, 7, 9.

The nucleic acids of the invention also include nucleic acids having sequence  
5 similarity or sequence identity. Nucleic acids having sequence similarity are detected by hybridization under low stringency conditions, for example, at 50°C and 10XSSC (0.9 M NaCl/0.09 M sodium citrate) and remain bound when subjected to washing at 55°C in 1XSSC. Sequence identity can be determined by hybridization under stringent conditions, for example, at 50°C or higher and 0.1XSSC (9 mM NaCl/0.9 mM sodium citrate).  
10 Hybridization methods and conditions are well known in the art, see U.S. Patent No. 5,707,829. Nucleic acids that are substantially identical to the provided nucleic acid sequences, e.g. allelic variants, genetically altered versions of the gene, etc., bind to the provided nucleic acid sequences (SEQ ID NO:1, 2, 3, 5, 7, 9) under stringent hybridization conditions. By using probes, particularly labeled probes of DNA sequences, one can isolate  
15 homologous or related genes. The source of homologous genes can be any species.

Preferably, hybridization is performed using at least 15 contiguous nucleotides of SEQ ID NO:1, 2, 3, 5, 7, 9. The probe will preferentially hybridize with a nucleic acid or mRNA comprising the complementary sequence, allowing the identification and retrieval of the nucleic acids of the biological material that uniquely hybridize to the selected probe.  
20 Probes of more than 15 nucleotides can be used, e.g. probes of from about 18 nucleotides to not more than about 100 nucleotides, but 15 nucleotides generally represents sufficient sequence for unique identification.

The nucleic acids of the invention also include naturally occurring variants of the nucleotide sequences, e.g. degenerate variants, allelic variants, etc. Variants of the nucleic  
25 acids of the invention are identified by hybridization of putative variants with nucleotide sequences disclosed herein, preferably by hybridization under stringent conditions. For example, by using appropriate wash conditions, variants of the nucleic acids of the invention can be identified where the allelic variant exhibits at most about 25-30% base pair mismatches relative to the selected nucleic acid probe. In general, allelic variants contain  
30 5-25% base pair mismatches, and can contain as little as even 2-5%, or 1-2% base pair mismatches, as well as a single base-pair mismatch.

The invention also encompasses homologs corresponding to the nucleic acids of SEQ ID NO:1, 2, 3, 5, 7, 9, where the source of homologous genes can be any related

species within the same genus or group. Within a group, homologs have substantial sequence similarity, e.g. at least 75% sequence identity, usually at least 90%, more usually at least 95% between nucleotide sequences. Sequence similarity is calculated based on a reference sequence, which may be a subset of a larger sequence, such as a conserved motif, coding region, flanking region, etc. A reference sequence will usually be at least about 18 contiguous nt long, more usually at least about 30 nt long, and may extend to the complete sequence that is being compared. Algorithms for sequence analysis are known in the art, such as BLAST, described in Altschul et al., J. Mol. Biol. (1990) 215:403-10.

In general, variants of the invention have a sequence identity greater than at least about 65%, preferably at least about 75%, more preferably at least about 85%, and can be greater than at least about 90% or more as determined by the Smith-Waterman homology search algorithm as implemented in MPSRCH program (Oxford Molecular). For the purposes of this invention, a preferred method of calculating percent identity is the Smith-Waterman algorithm, using the following. Global DNA sequence identity must be greater than 65% as determined by the Smith-Waterman homology search algorithm as implemented in MPSRCH program (Oxford Molecular) using an affine gap search with the following search parameters: gap open penalty, 12; and gap extension penalty, 1.

ATnov polymorphic sequences. It has been found that specific sites in the ATnov gene sequence are polymorphic, i.e. within a population, more than one nucleotide (G, A, T, C) is found at a specific position. Polymorphisms may provide functional differences in the genetic sequence, through changes in the encoded polypeptide, changes in mRNA stability, binding of transcriptional and translation factors to the DNA or RNA, and the like.

The polymorphisms are also used as single nucleotide polymorphisms to detect association with, or genetic linkage to phenotypic variation in activity and expression of ATnov.

SNPs are generally biallelic systems, that is, there are two alleles that an individual may have for any particular marker. SNPs, found approximately every kilobase, offer the potential for generating very high density genetic maps, which will be extremely useful for developing haplotyping systems for genes or regions of interest, and because of the nature of SNPs, they may in fact be the polymorphisms associated with the disease phenotypes under study. The low mutation rate of SNPs also makes them excellent markers for studying complex genetic traits.



Single nucleotide polymorphisms are provided in the ATnov3 sequence listing. The provided sequences also encompass the complementary sequence corresponding to any of the provided polymorphisms.

5 In order to provide an unambiguous identification of the specific site of a polymorphism, sequences flanking the polymorphic site are included in a probe for the region. It will be understood that there is no special significance to the length of non-polymorphic flanking sequence that is included, except to aid in positioning the polymorphism in the genomic sequence.

10 For screening purposes, hybridization probes of the polymorphic sequences may be used where both forms are present, either in separate reactions, spatially separated on a solid phase matrix, or labeled such that they can be distinguished from each other. Assays may utilize nucleic acids that hybridize to one or more of the described polymorphisms.

15 An array may include all or a subset of the ATnov3 polymorphisms. One or both polymorphic forms may be present in the array. Usually such an array will include at least 2 different polymorphic sequences, i.e. polymorphisms located at unique positions within the locus, and may include as many all of the provided polymorphisms. Arrays of interest may further comprise sequences, including polymorphisms, of other genetic sequences, particularly other sequences of interest for pharmacogenetic screening. The oligonucleotide sequence on the array will usually be at least about 12 nt in length, may be the length of the  
20 provided polymorphic sequences, or may extend into the flanking regions to generate fragments of 100 to 200 nt in length. For examples of arrays, see Ramsay (1998) Nat. Biotech. 16:40-44; Hacia et al. (1996) Nature Genetics 14:441-447; Lockhart et al. (1996) Nature Biotechnol. 14:1675-1680; and De Risi et al. (1996) Nature Genetics 14:457-460.

25 The subject nucleic acids can be cDNAs or genomic DNAs, as well as fragments thereof, particularly fragments that encode a biologically active gene product and/or are useful in the methods disclosed herein. The term "cDNA" as used herein is intended to include all nucleic acids that share the arrangement of sequence elements found in native mature mRNA species, where sequence elements are exons and 3' and 5' non-coding regions. Normally mRNA species have contiguous exons, with the intervening introns, when  
30 present, being removed by nuclear RNA splicing, to create a continuous open reading frame encoding a polypeptide of the invention.

A genomic sequence of interest comprises the nucleic acid present between the initiation codon and the stop codon, as defined in the listed sequences, including all of the

introns that are normally present in a native chromosome. It can further include the 3' and 5' untranslated regions found in the mature mRNA. It can further include specific transcriptional and translational regulatory sequences, such as promoters, enhancers, etc., including about 1 kb, but possibly more, of flanking genomic DNA at either the 5' and 3' end of the transcribed region. The genomic DNA can be isolated as a fragment of 100 kbp or smaller; and substantially free of flanking chromosomal sequence. The genomic DNA flanking the coding region, either 3' and 5', or internal regulatory sequences as sometimes found in introns, contains sequences required for expression.

The nucleic acid compositions of the subject invention can encode all or a part of the subject polypeptides. Double or single stranded fragments can be obtained from the DNA sequence by chemically synthesizing oligonucleotides in accordance with conventional methods, by restriction enzyme digestion, by PCR amplification, etc. Isolated nucleic acids and nucleic acid fragments of the invention comprise at least about 15 up to about 100 contiguous nucleotides, or up to the complete sequence provided in SEQ ID NO:1, 2, 3, 5, 7 or 9. For the most part, fragments will be of at least 15 nt, usually at least 18 nt or 25 nt, and up to at least about 50 contiguous nt in length or more.

Probes specific to the nucleic acids of the invention can be generated using the nucleic acid sequences disclosed in SEQ ID NO:1, 2, 3, 5, 7 or 9 and the fragments as described above. The probes can be synthesized chemically or can be generated from longer nucleic acids using restriction enzymes. The probes can be labeled, for example, with a radioactive, biotinylated, or fluorescent tag. Preferably, probes are designed based upon an identifying sequence of a nucleic acid of one of SEQ ID NO:1, 2, 3, 5, 7 or 9. More preferably, probes are designed based on a contiguous sequence of one of the subject nucleic acids that remain unmasked following application of a masking program for masking low complexity (e.g., XBLAST) to the sequence., i.e. one would select an unmasked region, as indicated by the nucleic acids outside the poly-n stretches of the masked sequence produced by the masking program.

The nucleic acids of the subject invention are isolated and obtained in substantial purity, generally as other than an intact chromosome. Usually, the nucleic acids, either as DNA or RNA, will be obtained substantially free of other naturally-occurring nucleic acid sequences, generally being at least about 50%, usually at least about 90% pure and are typically "recombinant", e.g., flanked by one or more nucleotides with which it is not normally associated on a naturally occurring chromosome.

The nucleic acids of the invention can be provided as a linear molecule or within a circular molecule. They can be provided within autonomously replicating molecules (vectors) or within molecules without replication sequences. They can be regulated by their own or by other regulatory sequences, as is known in the art. The nucleic acids of the invention can be introduced into suitable host cells using a variety of techniques which are available in the art, such as transferrin polycation-mediated DNA transfer, transfection with naked or encapsulated nucleic acids, liposome-mediated DNA transfer, intracellular transportation of DNA-coated latex beads, protoplast fusion, viral infection, electroporation, gene gun, calcium phosphate-mediated transfection, and the like.

The subject nucleic acid compositions can be used to, for example, produce polypeptides, as probes for the detection of mRNA of the invention in biological samples (e.g., extracts of cells) to generate additional copies of the nucleic acids, to generate ribozymes or antisense oligonucleotides, and as single stranded DNA probes or as triple-strand forming oligonucleotides. The probes described herein can be used to, for example, determine the presence or absence of the nucleic acid sequences as shown in SEQ ID NO:1, 2, 3, 5, 7 or 9 or variants thereof in a sample.

The sequence of the 5' flanking region may be utilized for promoter elements, including enhancer binding sites, that provide for developmental regulation in tissues where ATnov genes are expressed. The tissue specific expression is useful for determining the pattern of expression, and for providing promoters that mimic the native pattern of expression. Naturally occurring polymorphisms in the promoter regions are useful for determining natural variations in expression, particularly those that may be associated with disease.

Alternatively, mutations may be introduced into the promoter regions to determine the effect of altering expression in experimentally defined systems. Methods for the identification of specific DNA motifs involved in the binding of transcriptional factors are known in the art, e.g. sequence similarity to known binding motifs, gel retardation studies, etc. For examples, see Blackwell et al. (1995) Mol Med 1: 194-205; Mortlock et al. (1996) Genome Res. 6: 327-33; and Joulin and Richard-Foy (1995) Eur J Biochem 232: 620-626.

The regulatory sequences may be used to identify cis acting sequences required for transcriptional or translational regulation of ATnov expression, especially in different tissues or stages of development, and to identify cis acting sequences and trans acting factors that regulate or mediate ATnov expression. Such transcription or translational control regions

may be operably linked to a ATnov gene in order to promote expression of wild type or altered ATnov or other proteins of interest in cultured cells, or in embryonic, fetal or adult tissues, and for gene therapy.

Double or single stranded fragments may be obtained of the DNA sequence by  
5 chemically synthesizing oligonucleotides in accordance with conventional methods, by restriction enzyme digestion, by PCR amplification, etc. For the most part, DNA fragments will be of at least 15 nt, usually at least 18 nt or 25 nt, and may be at least about 50 nt. Such small DNA fragments are useful as primers for PCR, hybridization screening probes, etc. Larger DNA fragments, i.e. greater than 100 nt are useful for production of the encoded  
10 polypeptide. For use in amplification reactions, such as PCR, a pair of primers will be used. The exact composition of the primer sequences is not critical to the invention, but for most applications the primers will hybridize to the subject sequence under stringent conditions, as known in the art. It is preferable to choose a pair of primers that will generate an amplification product of at least about 50 nt, preferably at least about 100 nt. Algorithms for  
15 the selection of primer sequences are generally known, and are available in commercial software packages. Amplification primers hybridize to complementary strands of DNA, and will prime towards each other.

The DNA may also be used to identify expression of the gene in a biological specimen. The manner in which one probes cells for the presence of particular nucleotide  
20 sequences, as genomic DNA or RNA, is well established in the literature and does not require elaboration here. DNA or mRNA is isolated from a cell sample. The mRNA may be amplified by RT-PCR, using reverse transcriptase to form a complementary DNA strand, followed by polymerase chain reaction amplification using primers specific for the subject DNA sequences. Alternatively, the mRNA sample is separated by gel electrophoresis,  
25 transferred to a suitable support, e.g. nitrocellulose, nylon, etc., and then probed with a fragment of the subject DNA as a probe. Other techniques, such as oligonucleotide ligation assays, in situ hybridizations, and hybridization to DNA probes arrayed on a solid chip may also find use. Detection of mRNA hybridizing to the subject sequence is indicative of ATnov gene expression in the sample.

30 The sequence of an ATnov gene, including flanking promoter regions and coding regions, may be mutated in various ways known in the art to generate targeted changes in promoter strength, sequence of the encoded protein, etc. The DNA sequence or protein product of such a mutation will usually be substantially similar to the sequences provided

herein, i.e. will differ by at least one nucleotide or amino acid, respectively, and may differ by at least two but not more than about ten nucleotides or amino acids. The sequence changes may be substitutions, insertions or deletions. Deletions may further include larger changes, such as deletions of a domain or exon. Other modifications of interest include  
5 epitope tagging, e.g. with the FLAG system, HA, etc. For studies of subcellular localization, fusion proteins with green fluorescent proteins (GFP) may be used.

Techniques for in vitro mutagenesis of cloned genes are known. Examples of protocols for site specific mutagenesis may be found in Gustin et al., *Biotechniques* 14:22 (1993); Barany, *Gene* 37:111-23 (1985); Colicelli et al., *Mol Gen Genet* 199:537-9 (1985);  
10 and Prentki et al., *Gene* 29:303-13 (1984). Methods for site specific mutagenesis can be found in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, CSH Press 1989, pp. 15.3-15.108; Weiner et al., *Gene* 126:35-41 (1993); Sayers et al., *Biotechniques* 13:592-6 (1992); Jones and Winistorfer, *Biotechniques* 12:528-30 (1992); Barton et al., *Nucleic Acids Res* 18:7349-55 (1990); Marotti and Tomich, *Gene Anal Tech* 6:67-70 (1989); and Zhu, *Anal*  
15 *Biochem* 177:120-4 (1989). Such mutated genes may be used to study structure-function relationships of ATnov polypeptides, or to alter properties of the protein that affect its function or regulation.

Genetic polymorphisms, either naturally occurring or introduced as described above, are useful in screening for altered transport or metabolism of ATnov substrates. For  
20 example, variant alleles may affect the pharmacokinetic parameters of substrates. A drug's volume of distribution, clearance, and the derived parameter, half-life, are particularly important, as they determine the degree of fluctuation between a maximum and minimum plasma concentration during a dosage interval, the magnitude of steady state concentration and the time to reach steady state plasma concentration upon chronic dosing. Parameters  
25 derived from in vivo drug administration are useful in determining the clinical effect of a particular ATnov genotype.

#### ATNOV POLYPEPTIDES

The subject gene may be employed for producing all or portions of ATnov  
30 polypeptides. Fragments of interest include the glycosylation sites, transmembrane domains, ATP binding regions, the substrate binding sites, etc. Such domains will usually include at least about 20 amino acids of the provided sequence, more usually at least about 50 amino acids, and may include 100 amino acids or more, up to the complete domain.

Binding contacts may be comprised of non-contiguous sequences, which are brought into proximity by the tertiary structure of the protein. The sequence of such fragments may be modified through manipulation of the coding sequence, as described above. Truncations may be performed at the carboxy or amino terminus of the fragment, e.g. to determine the minimum sequence required for biological activity.

A subset of the provided nucleic acid polymorphisms in ATnov3 confer a change in the corresponding amino acid sequence, as previously described. Using the amino acid sequence provided in SEQ ID NO:3 as a reference, the amino acid polymorphisms of the invention include asndasp, pos. 130; and a frameshift at position 537 resulting in a truncated protein of 542 amino acids. Polypeptides comprising at least one of the provided polymorphisms (ATnov3' polypeptides) are of interest. The term "ATnov3' polypeptides" as used herein includes complete ATnov protein forms, e.g. such splicing variants as known in the art, and fragments thereof, which fragments may comprise short polypeptides, epitopes, functional domains; binding sites; etc.; and including fusions of the subject polypeptides to other proteins or parts thereof. Polypeptides will usually be at least about 8 amino acids in length, more usually at least about 12 amino acids in length, and may be 20 amino acids or longer, up to substantially the complete protein.

For expression, an expression cassette may be employed. The expression vector will provide a transcriptional and translational initiation region, which may be inducible or constitutive, where the coding region is operably linked under the transcriptional control of the transcriptional initiation region, and a transcriptional and translational termination region. These control regions may be native to an ATnov gene, or may be derived from exogenous sources.

The peptide may be expressed in prokaryotes or eukaryotes in accordance with conventional ways, depending upon the purpose for expression. For large scale production of the protein, a unicellular organism, such as *E. coli*, *B. subtilis*, *S. cerevisiae*, insect cells in combination with baculovirus vectors, or cells of a higher organism such as vertebrates, particularly mammals, e.g. COS 7 cells, may be used as the expression host cells. In some situations, it is desirable to express the ATnov gene in eukaryotic cells, where the ATnov protein will benefit from native folding and post-translational modifications. Small peptides can also be synthesized in the laboratory. Peptides that are subsets of the complete ATnov sequence may be used to identify and investigate parts of the protein important for function, or to raise antibodies directed against these regions.

With the availability of the protein or fragments thereof in large amounts, by employing an expression host, the protein may be isolated and purified in accordance with conventional ways. A lysate may be prepared of the expression host and the lysate purified using HPLC, exclusion chromatography, gel electrophoresis, affinity chromatography, or other purification technique. The purified protein will generally be at least about 80% pure, preferably at least about 90% pure, and may be up to and including 100% pure. Pure is intended to mean free of other proteins, as well as cellular debris.

The expressed ATnov polypeptides are useful for the production of antibodies, where short fragments provide for antibodies specific for the particular polypeptide, and larger fragments or the entire protein allow for the production of antibodies over the surface of the polypeptide. Antibodies may be raised to the wild-type or variant forms of ATnov. Antibodies may be raised to isolated peptides corresponding to these domains, or to the native protein.

Antibodies are prepared in accordance with conventional ways, where the expressed polypeptide or protein is used as an immunogen, by itself or conjugated to known immunogenic carriers, e.g. KLH, pre-S HBsAg, other viral or eukaryotic proteins, or the like. Various adjuvants may be employed, with a series of injections, as appropriate. For monoclonal antibodies, after one or more booster injections, the spleen is isolated, the lymphocytes immortalized by cell fusion, and then screened for high affinity antibody binding. The immortalized cells, i.e. hybridomas, producing the desired antibodies may then be expanded. For further description, see Monoclonal Antibodies: A Laboratory Manual, Harlow and Lane eds., Cold Spring Harbor Laboratories, Cold Spring Harbor, New York, 1988. If desired, the mRNA encoding the heavy and light chains may be isolated and mutagenized by cloning in *E. coli*, and the heavy and light chains mixed to further enhance the affinity of the antibody. Alternatives to in vivo immunization as a method of raising antibodies include binding to phage "display" libraries, usually in conjunction with in vitro affinity maturation.

#### ATNOV GENOTYPING

The subject nucleic acid and/or polypeptide compositions may be used in genotyping and to screen for the presence of polymorphisms in the sequence, or variation in the expression of the subject genes. Genotyping may be performed to determine whether a particular polymorphisms is associated with a disease state or genetic predisposition to a disease state, particularly diseases associated with liver disorders.

Genotyping may also be performed for pharmacogenetic analysis to assess the association between an individual's genotype and that individual's ability to react to a therapeutic agent. Differences in substrate transport to relevant cells can lead to toxicity or therapeutic failure. Relationships between polymorphisms in transporter expression or specificity can be used to optimize therapeutic dose administration.

ATnov genotyping is performed by DNA or RNA sequence and/or hybridization analysis of any convenient sample from a patient, e.g. biopsy material, blood sample, scrapings from cheek, etc. A nucleic acid sample from an individual is analyzed for the presence of polymorphisms in ATnov, particularly those that affect the activity, responsiveness or expression of ATnov. Specific sequences of interest include any polymorphism that leads to changes in basal expression in one or more tissues, to changes in the modulation of ATnov expression, or alterations in ATnov specificity and/or activity.

The effect of a polymorphism in ATnov gene sequence on the response to a particular agent may be determined by in vitro or in vivo assays. Such assays may include monitoring during clinical trials, testing on genetically defined cell lines, etc. The response of an individual to the agent can then be predicted by determining the ATnov genotype with respect to the polymorphism. Where there is a differential distribution of a polymorphism by racial background, guidelines for drug administration can be generally tailored to a particular ethnic group.

Biochemical studies may be performed to determine whether a sequence polymorphism in a ATnov coding region or control regions is associated with disease. Disease associated polymorphisms may include deletion or truncation of the gene, mutations that alter expression level, that affect the specificity or transport kinetics of the transporter, etc.

A number of methods are available for analyzing nucleic acids for the presence of a specific sequence. Where large amounts of DNA are available, genomic DNA is used directly. Alternatively, the region of interest is cloned into a suitable vector and grown in sufficient quantity for analysis. The nucleic acid may be amplified by conventional techniques, such as the polymerase chain reaction (PCR), to provide sufficient amounts for analysis. The use of the polymerase chain reaction is described in Saiki et al. (1985) Science 239:487, and a review of current techniques may be found in Sambrook et al. Molecular Cloning: A Laboratory Manual, CSH Press 1989, pp.14.2-14.33. Amplification may be used to determine whether a polymorphism is present, by using a primer that is



specific for the polymorphism. Alternatively, various methods are known in the art that utilize oligonucleotide ligation as a means of detecting polymorphisms, for examples see Delahunty et al. (1996) Am. J. Hum. Genet. 58:1239-1246.

A detectable label may be included in an amplification reaction. Suitable labels  
5 include fluorochromes, e.g. fluorescein isothiocyanate (FITC), rhodamine, Texas Red, phycoerythrin, allophycocyanin, 6-carboxyfluorescein (6-FAM), 2',7'-dimethoxy-4',5'-dichloro-6-carboxyfluorescein (JOE), 6-carboxy-X-rhodamine (ROX), 6-carboxy-2',4',7',4,7-hexachlorofluorescein (HEX), 5-carboxyfluorescein (5-FAM) or N,N,N',N'-tetramethyl-6-carboxyrhodamine (TAMRA), radioactive labels, e.g.  $^{32}\text{P}$ ,  $^{35}\text{S}$ ,  $^3\text{H}$ ; etc. The label may be a  
10 two stage system, where the amplified DNA is conjugated to biotin, haptens, etc. having a high affinity binding partner, e.g. avidin, specific antibodies, etc., where the binding partner is conjugated to a detectable label. The label may be conjugated to one or both of the primers. Alternatively, the pool of nucleotides used in the amplification is labeled, so as to incorporate the label into the amplification product.

15 The sample nucleic acid, e.g. amplified or cloned fragment, is analyzed by one of a number of methods known in the art. The nucleic acid may be sequenced by dideoxy or other methods. Hybridization with the variant sequence may also be used to determine its presence, by Southern blots, dot blots, etc. The hybridization pattern of a control and variant sequence to an array of oligonucleotide probes immobilised on a solid support, as described  
20 in U.S. 5,445,934, or in WO95/35505, may also be used as a means of detecting the presence of variant sequences. Single strand conformational polymorphism (SSCP) analysis, denaturing gradient gel electrophoresis (DGGE), mismatch cleavage detection, and heteroduplex analysis in gel matrices are used to detect conformational changes created by DNA sequence variation as alterations in electrophoretic mobility. Alternatively, where a  
25 polymorphism creates or destroys a recognition site for a restriction endonuclease (restriction fragment length polymorphism, RFLP), the sample is digested with that endonuclease, and the products size fractionated to determine whether the fragment was digested. Fractionation is performed by gel or capillary electrophoresis, particularly acrylamide or agarose gels.

30 In one embodiment of the invention, an array of oligonucleotides are provided, where discrete positions on the array are complementary to one or more of the provided polymorphic sequences, e.g. oligonucleotides of at least 12 nt, frequently 20 nt, or larger, and including the sequence flanking the polymorphic position. Such an array may comprise

a series of oligonucleotides, each of which can specifically hybridize to a different polymorphism. For examples of arrays, see Hacia et al. (1996) Nature Genetics 14:441-447; Lockhart et al. (1996) Nature Biotechnol. 14:1675-1680; and De Risi et al. (1996) Nature Genetics 14:457-460.

5           Screening for polymorphisms in ATnov may be based on the functional or antigenic characteristics of the protein. Protein truncation assays are useful in detecting deletions that may affect the biological activity of the protein. Various immunoassays designed to detect polymorphisms in ATnov proteins may be used in screening. Where many diverse genetic mutations lead to a particular disease phenotype, functional protein assays have proven to  
10 be effective screening tools. The activity of the encoded ATnov protein as a anion transporter may be determined by comparison with the wild-type protein.

          Antibodies specific for a ATnov may be used in staining or in immunoassays. Samples, as used herein, include biological fluids such as semen, blood, cerebrospinal fluid, tears, saliva, lymph, dialysis fluid and the like; organ or tissue culture derived fluids; and  
15 fluids extracted from physiological tissues. Also included in the term are derivatives and fractions of such fluids. The cells may be dissociated, in the case of solid tissues, or tissue sections may be analyzed. Alternatively a lysate of the cells may be prepared.

          Diagnosis may be performed by a number of methods to determine the absence or presence or altered amounts of normal or abnormal ATnov polypeptides in patient cells. For  
20 example, detection may utilize staining of cells or histological sections, performed in accordance with conventional methods. The antibodies of interest are added to the cell sample, and incubated for a period of time sufficient to allow binding to the epitope, usually at least about 10 minutes. The antibody may be labeled with radioisotopes, enzymes, fluorsceners, chemiluminescers, or other labels for direct detection. Alternatively, a second  
25 stage antibody or reagent is used to amplify the signal. Such reagents are well known in the art. For example, the primary antibody may be conjugated to biotin, with horseradish peroxidase-conjugated avidin added as a second stage reagent. Alternatively, the secondary antibody conjugated to a flourescent compound, e.g. flourescein, rhodamine, Texas red, etc. Final detection uses a substrate that undergoes a color change in the  
30 presence of the peroxidase. The absence or presence of antibody binding may be determined by various methods, including flow cytometry of dissociated cells, microscopy, radiography, scintillation counting, etc.

## MODULATION OF GENE EXPRESSION

The ATnov genes, gene fragments, or the encoded protein or protein fragments are useful in gene therapy to treat disorders associated with ATnov defects. Expression vectors may be used to introduce the ATnov gene into a cell. Such vectors generally have  
5 convenient restriction sites located near the promoter sequence to provide for the insertion of nucleic acid sequences. Transcription cassettes may be prepared comprising a transcription initiation region, the target gene or fragment thereof, and a transcriptional termination region. The transcription cassettes may be introduced into a variety of vectors, e.g. plasmid; retrovirus, e.g. lentivirus; adenovirus; and the like, where the vectors are able  
10 to transiently or stably be maintained in the cells, usually for a period of at least about one day, more usually for a period of at least about several days to several weeks.

The gene or ATnov protein may be introduced into tissues or host cells by any number of routes, including viral infection, microinjection, or fusion of vesicles. Jet injection may also be used for intramuscular administration, as described by Furth et al. (1992) Anal  
15 Biochem 205:365-368. The DNA may be coated onto gold microparticles, and delivered intradermally by a particle bombardment device, or "gene gun" as described in the literature (see, for example, Tang et al. (1992) Nature 356:152-154), where gold microprojectiles are coated with the ATnov or DNA, then bombarded into skin cells.

Antisense molecules can be used to down-regulate expression of ATnov in cells.  
20 The anti-sense reagent may be antisense oligonucleotides (ODN), particularly synthetic ODN having chemical modifications from native nucleic acids, or nucleic acid constructs that express such anti-sense molecules as RNA. The antisense sequence is complementary to the mRNA of the targeted gene, and inhibits expression of the targeted gene products. Antisense molecules inhibit gene expression through various mechanisms, e.g. by reducing  
25 the amount of mRNA available for translation, through activation of RNase H, or steric hindrance. One or a combination of antisense molecules may be administered, where a combination may comprise multiple different sequences.

Antisense molecules may be produced by expression of all or a part of the target gene sequence in an appropriate vector, where the transcriptional initiation is oriented such  
30 that an antisense strand is produced as an RNA molecule. Alternatively, the antisense molecule is a synthetic oligonucleotide. Antisense oligonucleotides will generally be at least about 7, usually at least about 12, more usually at least about 20 nucleotides in length, and not more than about 500, usually not more than about 50, more usually not more than about

35 nucleotides in length, where the length is governed by efficiency of inhibition, specificity, including absence of cross-reactivity, and the like. It has been found that short oligonucleotides, of from 7 to 8 bases in length, can be strong and selective inhibitors of gene expression (see Wagner et al. (1996) Nature Biotechnology 14:840-844).

5 A specific region or regions of the endogenous sense strand mRNA sequence is chosen to be complemented by the antisense sequence. Selection of a specific sequence for the oligonucleotide may use an empirical method, where several candidate sequences are assayed for inhibition of expression of the target gene in an in vitro or animal model. A combination of sequences may also be used, where several regions of the mRNA sequence  
10 are selected for antisense complementation.

Antisense oligonucleotides may be chemically synthesized by methods known in the art (see Wagner et al. (1993) *supra*. and Milligan et al., *supra*.) Preferred oligonucleotides are chemically modified from the native phosphodiester structure, in order to increase their intracellular stability and binding affinity. A number of such modifications have been  
15 described in the literature, which alter the chemistry of the backbone, sugars or heterocyclic bases.

Among useful changes in the backbone chemistry are phosphorothioates; phosphorodithioates, where both of the non-bridging oxygens are substituted with sulfur; phosphoroamidites; alkyl phosphotriesters and boranophosphates. Achiral phosphate  
20 derivatives include 3'-O-5'-S-phosphorothioate, 3'-S-5'-O-phosphorothioate, 3'-CH<sub>2</sub>-5'-O-phosphonate and 3'-NH-5'-O-phosphoroamidate. Peptide nucleic acids replace the entire ribose phosphodiester backbone with a peptide linkage. Sugar modifications are also used to enhance stability and affinity. The  $\alpha$ -anomer of deoxyribose may be used, where the base is inverted with respect to the natural  $\beta$ -anomer. The 2'-OH of the ribose sugar  
25 may be altered to form 2'-O-methyl or 2'-O-allyl sugars, which provides resistance to degradation without comprising affinity. Modification of the heterocyclic bases must maintain proper base pairing. Some useful substitutions include deoxyuridine for deoxythymidine; 5-methyl-2'-deoxycytidine and 5-bromo-2'-deoxycytidine for deoxycytidine. 5-propynyl-2'-deoxyuridine and 5-propynyl-2'-deoxycytidine have been shown to increase affinity and  
30 biological activity when substituted for deoxythymidine and deoxycytidine, respectively.

As an alternative to anti-sense inhibitors, catalytic nucleic acid compounds, e.g. ribozymes, anti-sense conjugates, etc. may be used to inhibit gene expression. Ribozymes may be synthesized in vitro and administered to the patient, or may be encoded on an

expression vector, from which the ribozyme is synthesized in the targeted cell (for example, see International patent application WO 9523225, and Beigelman et al. (1995) Nucl. Acids Res 23:4434-42). Examples of oligonucleotides with catalytic activity are described in WO 9506764. Conjugates of anti-sense ODN with a metal complex, e.g. terpyridylCu(II), capable  
5 of mediating mRNA hydrolysis are described in Bashkin et al. (1995) Appl Biochem Biotechnol 54:43-56.

#### GENETICALLY ALTERED CELL OR ANIMAL MODELS FOR ATNOV FUNCTION

The subject nucleic acids can be used to generate transgenic animals or site specific  
10 gene modifications in cell lines. Transgenic animals may be made through homologous recombination, where the normal ATnov locus is altered. Alternatively, a nucleic acid construct is randomly integrated into the genome. Vectors for stable integration include plasmids, retroviruses and other animal viruses, YACs, and the like.

The modified cells or animals are useful in the study of ATnov function and  
15 regulation. For example, a series of small deletions and/or substitutions may be made in the ATnov gene to determine the role of different transmembrane domains, of ATP catalysis, etc. Of interest are the use of ATnov to construct transgenic animal models where expression of ATnov is specifically reduced or absent. Specific constructs of interest include anti-sense ATnov, which will block ATnov expression, expression of dominant negative ATnov  
20 mutations, etc. One may also provide for expression of the ATnov gene or variants thereof in cells or tissues where it is not normally expressed or at abnormal times of development.

DNA constructs for homologous recombination will comprise at least a portion of the ATnov gene with the desired genetic modification, and will include regions of homology to  
25 the target locus. DNA constructs for random integration need not include regions of homology to mediate recombination. Conveniently, markers for positive and negative selection are included. Methods for generating cells having targeted gene modifications through homologous recombination are known in the art. For various techniques for transfecting mammalian cells, see Keown et al. (1990) Methods in Enzymology 185:527-537.

30 For embryonic stem (ES) cells, an ES cell line may be employed, or embryonic cells may be obtained freshly from a host, e.g. mouse, rat, guinea pig, etc. Such cells are grown on an appropriate fibroblast-feeder layer or grown in the presence of leukemia inhibiting factor (LIF). When ES or embryonic cells have been transformed, they may be used to

produce transgenic animals. After transformation, the cells are plated onto a feeder layer in an appropriate medium. Cells containing the construct may be detected by employing a selective medium. After sufficient time for colonies to grow, they are picked and analyzed for the occurrence of homologous recombination or integration of the construct. Those colonies that are positive may then be used for embryo manipulation and blastocyst injection. Blastocysts are obtained from 4 to 6 week old superovulated females. The ES cells are trypsinized, and the modified cells are injected into the blastocoel of the blastocyst. After injection, the blastocysts are returned to each uterine horn of pseudopregnant females. Females are then allowed to go to term and the resulting offspring screened for the construct. By providing for a different phenotype of the blastocyst and the genetically modified cells, chimeric progeny can be readily detected.

The chimeric animals are screened for the presence of the modified gene and males and females having the modification are mated to produce homozygous progeny. If the gene alterations cause lethality at some point in development, tissues or organs can be maintained as allogeneic or congenic grafts or transplants, or in in vitro culture. The transgenic animals may be any non-human mammal, such as laboratory animals, domestic animals, etc. The transgenic animals may be used in functional studies, drug screening, etc.

#### TESTING OF ATNOV FUNCTION and RESPONSES

Anion transporters such as ATnov polypeptides are involved in multiple biologically important processes. Pharmacological agents designed to affect only specific transporter subtypes are of particular interest. The subject polypeptides may be used to test the specificity of novel compounds, and of analogs and derivatives of compounds known to be substrates, or to act on anion transporters.

Drug screening may be performed using an in vitro model, a genetically altered cell or animal, or purified ATnov protein. One can identify ligands or substrates that bind to, modulate or mimic the action of ATnov. Drug screening identifies agents that provide a replacement for ATnov function in abnormal cells. Of particular interest are screening assays for agents that have a low toxicity for human cells. A wide variety of assays may be used for this purpose, including monitoring cellular excitation and conductance, labeled in vitro protein-protein binding assays, electrophoretic mobility shift assays, immunoassays for protein binding, and the like. The purified protein may also be used for determination of

three-dimensional crystal structure, which can be used for modeling intermolecular interactions.

The term "agent" as used herein describes any molecule, e.g. protein or pharmaceutical, with the capability of altering or mimicking the physiological function of  
5 ATnov polypeptide. Generally a plurality of assay mixtures are run in parallel with different agent concentrations to obtain a differential response to the various concentrations. Typically, one of these concentrations serves as a negative control, i.e. at zero concentration or below the level of detection.

Candidate agents encompass numerous chemical classes, though typically they are  
10 organic molecules, preferably small organic compounds having a molecular weight of more than 50 and less than about 2,500 daltons. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding, and typically include at least an amine, carbonyl, hydroxyl or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or  
15 heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules including peptides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs or combinations thereof.

Candidate agents are obtained from a wide variety of sources including libraries of  
20 synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds and biomolecules, including expression of randomized oligonucleotides and oligopeptides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural or synthetically produced libraries and compounds  
25 are readily modified through conventional chemical, physical and biochemical means, and may be used to produce combinatorial libraries. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification, etc. to produce structural analogs.

Where the screening assay is a binding assay, one or more of the molecules may be  
30 joined to a label, where the label can directly or indirectly provide a detectable signal. Various labels include radioisotopes, fluorescers, chemilumescers, enzymes, specific binding molecules, particles, e.g. magnetic particles, and the like. Specific binding molecules include pairs, such as biotin and streptavidin, digoxin and antidigoxin etc. For the specific

binding members, the complementary member would normally be labeled with a molecule that provides for detection, in accordance with known procedures.

5 A variety of other reagents may be included in the screening assay. These include reagents like salts, neutral proteins, e.g. albumin, detergents, etc. that are used to facilitate optimal protein-protein binding and/or reduce non-specific or background interactions. Reagents that improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, etc. may be used. The mixture of components are added in any order that provides for the requisite binding. Incubations are performed at any suitable temperature, typically between 4 and 40°C. Incubation periods are selected for optimum activity, but may also be optimized to facilitate rapid high-throughput screening. 10 Typically between 0.1 and 1 hours will be sufficient.

The compounds having the desired pharmacological activity may be administered in a physiologically acceptable carrier to a host in a variety of ways, orally, topically, parenterally e.g. subcutaneously, intraperitoneally, by viral infection, intravascularly, etc. 15 Depending upon the manner of introduction, the compounds may be formulated in a variety of ways. The concentration of therapeutically active compound in the formulation may vary from about 0.1-100 wt.%. The pharmaceutical compositions can be prepared in various forms, such as granules, tablets, pills, suppositories, capsules, suspensions, salves, lotions and the like. Pharmaceutical grade organic or inorganic carriers and/or diluents suitable for oral and topical use can be used to make up compositions containing the therapeutically- 20 active compounds. Diluents known to the art include aqueous media, vegetable and animal oils and fats. Stabilizing agents, wetting and emulsifying agents, salts for varying the osmotic pressure or buffers for securing an adequate pH value, and skin penetration enhancers can be used as auxiliary agents.



## EXPERIMENTAL

The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the subject invention, and are not intended to limit the scope of what is regarded as the invention. Efforts have been made to ensure accuracy with respect to the numbers used (e.g. amounts, temperature, concentrations, etc.) but some experimental errors and deviations should be allowed for. Unless otherwise indicated, parts are parts by weight, molecular weight is average molecular weight, temperature is in degrees centigrade; and pressure is at or near atmospheric.

## Example 1

Three novel members of the OATP gene family, which are expressed in liver tissue, were cloned. These genes were isolated using trapped exons obtained from large-scale exon trapping of chromosome 12. The three anion transporters reported here are 70-80% identical to each other over the predicted protein sequence, and are each 40% identical to the reported OATP protein sequence (Kullack-Ublick et al., (1995) Gastroenterology 109:1274-1282). The chromosomal location of these three anion transporters, along with the mapping of OATP, suggests this gene-family is clustered on 12p12.

## Materials and Methods

cDNA Isolation. cDNA clones were isolated using the GeneTrapper system (Gibco-BRL). PCR primers within the trapped exons were used to detect which plasmid cDNA libraries contained the gene of interest. Oligonucleotide probes were designed: (SEQ ID NO:7) C12B\_120: GGGGCTCTGATTGATACAACGTG ; (SEQ ID NO:8) C12C\_151: ACTGTGGCACACGTGGGTCATGTAGGACAT) and the process proceeded according to the supplied protocol. cDNA clones were sequenced on an ABI 377 according to standard methods. The Primer Island Transposition kit was used according to the supplied protocol. Sequences were analyzed, edited, and assembled using the Sequencher software (Gene Codes).

Radiation Hybrid Mapping. RH mapping was achieved using the Stanford G3 panel DNAs (Research Genetics). DNA was aliquoted into 96-well trays, dried, and resuspended in PCR buffer prior to PCR amplification. 20 µl PCR reactions with standard conditions, 2.5

mM MgCl<sub>2</sub>, Taq Gold, and an annealing temperature of 60°C (for ATnov1 and 2) or 55°C (for ATnov3) were used to detect expression. The assays were done in duplicate and results were scored and map positions determined via the RH server at Stanford University <<http://www-shgc.stanford.edu/RH/G3index.html>>.

5

RT-PCR. RT-PCR was utilized to characterize the expression pattern of the novel anion transporters. This approach used RNA from 30 different tissues to generate first strand cDNA. Total RNA was purchased (Clontech, Invitrogen) and used to synthesize first strand cDNA using M-MLV reverse transcriptase and the supplied buffer (Gibco-BRL). The 20 µl reaction contained 5 µg total RNA, 100 ng of random primers, 10 mM DTT, 0.5 mM each dNTP, and an RNase inhibitor (Gibco-BRL). Identical reactions were set up without reverse transcriptase to control for DNA contamination in the RNA samples. The synthesis reaction proceeded for 1 hour at 37°C followed by 10 minutes at 95°C. These cDNAs, along with control cDNA synthesis reactions without reverse transcriptase, were diluted 1:5 and 2 µl of each sample were arrayed into 96-well trays, dried, and resuspended in PCR buffer prior to PCR amplification. The cDNAs were tested with primers with defined expression patterns to verify the presence of amplifiable cDNA from each tissue. Gene-specific primers were used to amplify the cDNAs in 20 µl PCR reactions with standard conditions, 2.5 mM MgCl<sub>2</sub>, Taq Gold, and an appropriate annealing temperature.

This approach provides for relatively high-throughput analysis of gene expression in a large set of tissues in a cost-efficient manner and provides qualitative analysis of gene expression only. Modifications can be employed, such as the use of internal control primers, limited cycling parameters, and dilution series to convert this to a quantitative experiment.

#### 25 Primers for ATnov1

RH primers (SEQ ID NO:11) CTGCTGCCAACTAACATTGC  
(SEQ ID NO:12) CACACACTAACCATGCCTCT  
237 bp product

RT-PCR primers (SEQ ID NO:13) TCCAGTCATTGGCTTTGCAC  
(SEQ ID NO:14) AAGAACCAATAAAGCTGCTTACT  
413 bp product

#### 30 Primers for ATnov2

RH primers (SEQ ID NO:15) GTGTTTGCTAGCCACCTTGA

(SEQ ID NO:16) GGCAACACTTCCTCAAAGTG

196 bp product

RT-PCR primers (SEQ ID NO:17) GATGCTTTCCTCTGTGCAGT

(SEQ ID NO:18) CCTTCAAGCCGAAGAAGGCT

5

259 bp product

Primers for ATnov3

RH primers (SEQ ID NO:19) AGGAGTTCCTGGTCCTTTCA

(SEQ ID NO:20) CAAGCTAGACTTCAGGCCTT

10

137 bp product

RT-PCR primers (SEQ ID NO:21) GAGGAATTCTAGCTCCAATATATT

(SEQ ID NO:22) GTCCTACATGACCCACGTGTG

96 bp product

15 Results

cDNA Isolation. Large-scale exon trapping was completed across a chromosome 12 cosmid library. Approximately 2400 exons were sequenced and analyzed by BLAST algorithms to identify exons with potentially interesting homologies. Two different exons, C12B\_120 and C12C\_151 were identified that were 87% identical to each other and ~68% identical to a cloned organic anion transporter, OATP (Kullak-Ubrick et al. (1995) Gastroenterology 109: 1274-1282.), at the DNA level and ~78% similar at the amino acid level. Full-length cDNA clones were isolated using GeneTrapper (Gibco-BRL) from a liver cDNA library (Gibco-BRL). The resulting clones, the largest being up to 3.0 kb, were end-sequenced using vector primers. If the end sequences provided insufficient coverage of the cDNA clones, a transposon approach was used to complete the sequence of the cDNA clone.

The cDNA clones isolated with C12B\_120 yielded two different sequence contigs, ATnov1 and ATnov2, which were ~89% identical to each other. ATnov2 is identical to C12B\_120. cDNA clones isolated with C12C\_151 generated a sequence contig, ATnov3, that was ~86% identical to the first two contigs. Conceptual translations yielded predicted proteins of 688-704 amino acids in length. A multiple alignment of these three proteins is shown in figure 1. These genes also show significant homology to a human organic anion

transporter OATP (~40% identity, 60% similarity) and to a human prostaglandin transporter (~32% identity, 51% similarity) over the length of the predicted proteins.

Chromosomal Localization. The exon trapped products used in the cDNA screens  
5 were trapped from a chromosome 12 cosmid library, suggesting that at least ATnov2 and  
ATnov3 map to chromosome 12. OATP had been previously reported to map to  
chromosome 12 (Kullak-Ubrick et al., supra.) Radiation hybrid mapping was used to confirm  
the localization of these to chromosome 12, as well as to map them and ATnov1 to a specific  
region on the chromosome. The Stanford G3 panel showed linkage of all four of these  
10 genes to the marker GATA91H01, which is extrapolated to a cytogenetic location of 12p12.

Expression Analysis. OATP is expressed in multiple tissues, including brain, lung,  
liver, kidney, and testes (Kullak-Ubrick et al., supra.) RT-PCR was utilized to characterize  
the expression pattern of the novel anion transporters. This approach used RNA from 30  
15 different tissues to generate first strand cDNA. These cDNAs were arrayed, along with  
control cDNA synthesis reactions without reverse transcriptase, into 96-well trays, dried and  
stored until needed. This resource provides for relatively high-throughput analysis of gene  
expression in a large set of tissues in a cost-efficient manner. RT-PCR in this fashion allows  
for qualitative analysis of gene expression only.

20 PCR was performed on these plates with gene-specific primers for each of the ATnov  
genes. ATnov1 is expressed in fetal and adult liver; ATnov2 is expressed in adult liver and  
mammary gland; ATnov3 is expressed in fetal liver, adult liver, brain, adipose tissue, skin,  
and testes.

25 The predicted positions of transmembrane domains in the ATnov3 polypeptide are  
as follows:

	ATnov3
Transmembrane domain 1	29-45
Transmembrane domain 2	94-104
Transmembrane domain 3	169-189
Transmembrane domain 4	207-227
Transmembrane domain 5	259-279
Transmembrane domain 6	336-356

Transmembrane domain 7	376-396
Transmembrane domain 8	410-430
Transmembrane domain 9	481-501
Transmembrane domain 10	537-557
Transmembrane domain 11	581-601
Transmembrane domain 12	627-647

These novel members of the organic anion transporter family are expressed in the liver. Based on homology to another organic anion transporter, they are likely to be present on the basolateral surface of the hepatocytes and mediate the uptake of both xenobiotics and endogenous compounds for metabolism by the cytochrome p450s, glucuronosyl transferases, and other metabolic enzymes known to be present in the liver. The ATnov genes are all expressed in the liver, with ATnov 2 and 3 also being expressed in a limited number of other tissues. The RT-PCR approach described herein has a high level of sensitivity, with the ability to detect a control transcript diluted down to an expression level equivalent to a frequency of  $1/10^7$ .

The map positions of these anion transporters suggest that they lie adjacent to each other on the proximal short arm of chromosome 12. The anion transporters described herein are only ~89% identical to each other at the DNA level, suggesting that these genes arose via a recombination mechanism, but have since diverged sufficiently such that it is unlikely that these genes are polymorphic within a given population.

All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

## WHAT IS CLAIMED IS:

1. An isolated nucleic acid encoding a mammalian ATnov protein.
2. An isolated nucleic acid according to Claim 1, wherein said ATnov protein has  
5 the amino acid sequence of SEQ ID NO:4, 6, 8, or 10.
3. An isolated nucleic acid according to Claim 1, wherein said ATnov protein has  
an amino acid sequence that is substantially identical to the amino acid sequence of SEQ  
ID NO:4, 6, 8, or 10.  
10
4. An isolated nucleic acid according to Claim 1, comprising the nucleotide  
sequence as set forth in SEQ ID NO:1,2, 3, 5, 7, or 9.
5. An isolated nucleic acid that hybridizes under stringent conditions to the  
15 nucleic acid sequence of claim 4.
6. An expression cassette comprising a transcriptional initiation region functional  
in an expression host, a nucleic acid having a sequence of the isolated nucleic acid  
according to Claim 1 under the transcriptional regulation of said transcriptional initiation  
20 region, and a transcriptional termination region functional in said expression host.
7. A cell comprising an expression cassette according to Claim 6 as part of an  
extrachromosomal element or integrated into the genome of a host cell as a result of  
introduction of said expression cassette into said host cell, and the cellular progeny of said  
25 host cell.
8. A method for producing mammalian ATnov protein, said method comprising:  
growing a cell according to Claim 7, whereby said mammalian ATnov protein is  
expressed; and  
30 isolating said ATnov protein free of other proteins.
9. A purified polypeptide composition comprising at least 50 weight % of the  
protein present as a ATnov protein or a fragment thereof.

10. A monoclonal antibody binding specifically to an ATnov protein.
11. A non-human transgenic animal model for ATnov gene function wherein said  
5 transgenic animal comprises an introduced alteration in an ATnov gene.
12. The animal model of claim 11, wherein said animal is heterozygous for said introduced alteration.
- 10 13. The animal model of claim 12, wherein said animal is homozygous for said introduced alteration.
14. The animal model of claim 12, wherein said introduced alteration is a knockout of endogenous ATnov gene expression.
- 15 15. An isolated nucleic acid probe comprising an ATnov 3 sequence polymorphism, as part of other than a naturally occurring chromosome.
16. A nucleic acid probe according to Claim 15, wherein said probe is conjugated  
20 to a detectable marker.
17. An array of oligonucleotides comprising:  
two or more probes for detection of ATnov3 locus polymorphisms.

## SEQUENCE LISTING

<110> Miller, Andrew  
Buckler, Alan  
Laubert, Boris

<120> Human Anion Transporter Genes

<130> SEQ-23P

<150> 60/095,835

<151> 1998-08-07

<160> 23

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 2595

<212> DNA

<213> H. sapiens

<400> 1

agaaaaagga	tggacttggt	gcagttgctg	tagcattcaa	agtcaagggtg	atcatttcaa	60
accaagcatc	agcaacaatt	aaaaatattc	acttggtatc	tgtagttaa	taatggacca	120
acatcaacat	ttgaataaaa	cagcagagtc	agcatcttca	gagaaaaaga	aaacaagacg	180
ctgcaatgga	ttcaagatgt	tcttggcagc	cctgtcattc	agctatattg	ctaaagcact	240
agggtggaatc	attatgaaaa	tttccatcac	tcaaatagaa	aggagatttg	acatattcctc	300
ttctcttgct	ggtttaattg	atggaagctt	tgaattgga	aatttgcttg	tgattgtatt	360
tgtaagttaac	tttgatcta	aactacacag	accgaagtta	attggaattg	gttgtctcct	420
tatgggaact	ggaagtattt	tgacatcttt	accacatttc	ttcatgggat	attataggta	480
ttctaaagaa	acccatatta	atccatcaga	aaattcaaca	tcaagtttat	caacctgttt	540
aattaatcaa	accttatcat	tcaatggaac	atcacctgag	atagtagaaa	aagattgtgt	600
aaaggaatct	gggtcacaca	tgtggatcta	tgtcttcatg	gggaatatgc	ttcgtggcat	660
aggggaaacc	cccatagtac	cattggggat	ttcatacatt	gatgattttg	caaaagaagg	720
acattcttcc	ttgtatttag	gtagtttgaa	tgcaatagga	atgattgggc	cagtcattgg	780
ctttgcactg	ggatctctgt	ttgctaaaat	gtacgtggat	attggatatg	tagatctgag	840
cactatcaga	ataactccta	aggactctcg	ttgggttggg	gcttggtggc	ttggtttcct	900
tgtgtctgga	ctattttcca	ttatttcttc	cataccattt	tttttcttg	ccgaaaaatc	960
caaataaacc	acaaaaagaa	agaaaaattt	cactatcatt	gcatgtgctg	aaaacaaatg	1020
atgatagaaa	tcaaacagct	aatttgacca	accaaggaaa	aaatgttacc	aaaaatgtga	1080
ctgggttttt	ccagtctttg	aaaagcatcc	ttaccaatcc	cctgtatgtt	atatttctgc	1140
ttttgacatt	gttacaagta	agcagcttta	ttgggtcttt	tacttacgtc	tttaaataata	1200
tggagcaaca	gtacggtcag	tctgcatctc	atgctaactt	tttgttggga	atcataacca	1260
ttcctacggt	tgcaactgga	atgttttttag	gaggatttat	cattaaaaaa	ttcaaattgt	1320
ctttagttag	aattggcaaa	ttttcatttc	ttacttcgat	gatatccttc	ttgtttcaac	1380
ttctatattt	ccctctaata	tgcgaaagca	aatcagttgc	cggcctaacc	ttgacctatg	1440
atggaaataa	ttcagtggca	tctcatgtag	atgtaccact	ttcttattgc	aaactcagagt	1500
gcaattgtga	tgaaagtcag	tgggaaccag	tctgtgggaa	caatggaata	acttacctgt	1560
cacctgtgtc	agcaggatgc	aaatcctcaa	gtggtattaa	aaagcataca	gtgttttata	1620
actgtagtgt	tgtggaagta	actggtctcc	agaacagaaa	ttactcagca	cacttgggtg	1680
aatgcccaag	agataatact	tgtacaagga	aatttttcat	ctatgttgca	attcaagtca	1740
taaaactcttt	gttctctgca	acaggaggta	ccacatttat	cttgttgact	gtgaagattg	1800
ttcaacctga	attgaaagca	cttgcaatgg	gtttccagtc	aatggttata	agaacactag	1860
gaggaattct	agctccaata	tattttgggg	ctctgattga	taaaacatgt	atgaagtggg	1920
ccaccaacag	ctgtggagca	caaggggcct	gtaggatata	taattccgta	tttttgggaa	1980
gggtctactt	gggcttatct	atagctttta	gattcccagc	acttgtttta	tatattgttt	2040
tcatttttgc	tatgaagaaa	aaatttcaag	gaaaagatac	caaggcatcg	gacaatgaaa	2100
gaaaagtaat	ggatgaagca	aacttagaat	tcttaaataa	tggtgaacat	tttgtacctt	2160
ctgctggaac	agatagtaaa	acatgtaatt	tggacatgca	agacaatgct	gctgccaact	2220



aacattgcat	tgattcatta	agatgttatt	tttgaggtgt	tcctggtctt	tcactgacaa	2280
ttccaacatt	ctttacttac	agtggaccaa	tggataagtc	tatgcatcta	taataaacta	2340
taaaaaatgg	gagtacccat	ggttaggata	tagctatgcc	tttatggtta	agattagaat	2400
atatgatcca	taaaaaat	aagtggagag	catggttagt	gtgtgataca	ataaaaagta	2460
attgtttggt	agttgtaact	gctaataaaa	ccagtgacta	gaatataagg	gaggtaaaaa	2520
ggacaagata	gattaatagc	ctaaataaag	agaaaagcct	gatgccttta	aaaaaaaatg	2580
aaaaaaaaaa	aaaaa					2595

&lt;210&gt; 2

&lt;211&gt; 3273

&lt;212&gt; DNA

&lt;213&gt; H. sapiens

&lt;400&gt; 2

cccacgcgtc	cgatcagaaa	aaggatggac	ttgttgca	tgctgtagca	ttcaaagtca	60
aggtgatcat	ttcaaaccac	gcatcagcaa	caattaaaaa	tattcacttg	gtatctgtag	120
tttaataatg	gaccaacatc	aacatttgaa	taaaacagca	gagtcagcat	cttcagagaa	180
aaagaaaaca	agacgctgca	atggattcaa	gatgttcttg	gcagccctgt	cattcagcta	240
tattgctaaa	gcactagggtg	gaatcattat	gaaaatttcc	atcactcaaa	tagaaaggag	300
atttgacata	tcctcttctc	ttgctggttt	aattgatgga	agctttgaaa	ttggaaattt	360
gcttgtgatt	gtatttgtaa	gttactttgg	atctaaacta	cacagaccga	agttaattgg	420
aattggttgt	ctccttatgg	gaactggaag	tattttgaca	gctttaccac	atttcttcat	480
gggataatct	tcttgacact	cctgtcattc	agctatgttg	ctaaagcact	agctggaatt	540
tttatgaaaa	tatcaaccac	tcaaatagaa	aggagatttg	agatatcctc	ttctcttgtt	600
ggtttaattg	atggaagcct	cgaaatagga	aatttggttg	tgattgtatt	tgtaagttac	660
tttggatcta	aactacacag	accgaagtta	attggaattg	gttgttttct	tatgggaact	720
ggaagtattt	tgatggcttt	accacatttc	ttcatgggat	attacaggta	ttctaagaa	780
accaatattg	atccatcaga	aaattcaaca	tcaaacttac	caaactgttt	aattaatcaa	840
atgttatcac	tcaatagaac	accgtctgag	ataatagaaa	gaggttgtgt	gaaggaatct	900
gggtcacaca	tgtggatcta	tgtcttcatt	ggtaatatgc	ttcgtggcat	aggggaaacc	960
cccatagtac	cattggggat	ttcttacatt	gatgattttg	caaaagaagg	acattcttcc	1020
ttgtatttag	gtactgtgaa	tgcaatggga	atgactggtc	tagtttttgc	ctttatgctg	1080
ggatctctgt	ttgctaaaaa	gtatgtggat	atcggatatg	tggaactgag	cactatcagg	1140
ataactccta	aggactctcg	ttgggttggg	gcttgggtgc	ttggtttcct	tgtgtctgga	1200
atagtatcca	ttattttctc	tataccattc	tttttcttgc	ctctaaatcc	aaataaacca	1260
cagaaagaaa	ggaaagtctc	actatttttg	catgtgctaa	aaactaatga	taaaaggaat	1320
caaatagcta	atttgaccaa	ccgaagaaaa	tatattacca	aaaaatgtgac	tggtcttttc	1380
cagtctttga	aaagcatcct	taccaatccc	ctgtatgtta	tatttgaat	ttttacattg	1440
ttacacatga	gcagctacat	tgcttctctt	acttatatca	ttaaaatggg	ggagcaacag	1500
tatggttggt	ctgcatctaa	gactaacttt	ttgttgggag	tcctcgccct	acctactgtt	1560
gcaattggca	tgttttcagg	aggatatatc	attaaaaaat	tcaaattgtc	tttagttgga	1620
cttgccaaat	tggcattttg	ttctgcaaca	gtgcatctct	tatctcaagt	tttatatttc	1680
tttctaattc	gtgaaagcaa	atcagttggc	ggcctaacct	tgacctatga	tggaatatgt	1740
ccagtaagat	ctcatgtaga	tgtaccactt	tcttattgca	actcagagtg	caatttgtat	1800
gaaagtcaat	gggaaccggt	ctgtggaaac	aatggaataa	cttacctgtc	accttgtcta	1860
gcaggatgca	aatcttcaag	tgtaataaaa	gagcccatag	tggtttataa	ctgtagctgt	1920
gtggaagtta	ttggtctcca	gaacaaaaat	tactcagcgc	acttgggtga	atgcccaaga	1980
gatgatgctt	gtacaaggaa	atcttacggt	tattttgtaa	ttcaagtctt	agatgctttc	2040
ctctgtgcag	ttggacttac	ctcatatttc	gtgctggtga	ttaggattgt	tcaacctgaa	2100
ttgaaagcac	ttgcaatcgg	cttccattca	atgattatgc	gatcgctagg	aggtattcta	2160
gttccaatat	attttggggc	tctgattgat	acaacgtgta	tgaagtggtc	caccaacagc	2220
tgtggagcac	gaggggcttg	taggatatat	aattccacat	atttgggaag	agccttcttc	2280
ggcttgaagg	tagccctaata	atttccagta	cttgttttac	ttactgtatt	tatttttgtt	2340
gtaaggaaaa	aatcccatgg	aaaggatacc	aaagtattag	aaaatgaaag	acaagtaatt	2400
gatgaagcaa	acttagaatt	cttaaacgac	agtgaacatt	ttgtaccttc	tgctgaagaa	2460
cagtaaagca	tgtaattaa	aggaggaaaa	aaataatttt	gctgctgttt	ccaactaatg	2520
tattgattcc	ataagacgtt	atttttgtgg	tgctctgagt	cttttcaactg	agaattccca	2580
cattcttccac	ttatgatgca	acaatgaata	agcctatgaa	tttataatga	aacaaactat	2640
aaaaaatggt	acccatgggt	aggacatagc	tacacaagca	tttgtagttt	agaatatata	2700
attcataaaa	atttgaagtg	agaggaaatg	ttaatatgta	atagaagaaa	aagtactgtc	2760
tcaggtagtt	gtaactctta	ataaaaccaa	tgactagaat	acaagtggaa	gtaaaaaggt	2820
ggagatagat	taatagccta	aataacgaga	gaaccttatg	ccttttttaa	aacaaaacaa	2880

```

aaccattgag acattttact tagtcctaaa atctagcctg gatttatgct ataatgatat 2940
ctatttttca tgttaaattg tacattactc agaaattata aatattatta ctttataatt 3000
tgaaattgtg tttgctagcc accttgatgt attttcttcc aaactcccat taagatacta 3060
ttgaaaaaat agaaatagtc aaatatttgc aaggtataat tgtaggcaa catattatag 3120
catgtgttaa gtttctgcta ggcctatgga aatttttttt tttatttttg ttccattttt 3180
attcactttg aggaagtgtt gccttttttt ttgatgtact taaatggcta aaataaaaaa 3240
gacaatcaca agaaaaaaa aaaaaaaaaa aaa 3273

```

&lt;210&gt; 3

&lt;211&gt; 2799

&lt;212&gt; DNA

&lt;213&gt; H. sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (100)...(1729)

&lt;223&gt; ATnov3.1 Coding sequence

&lt;221&gt; variation

&lt;222&gt; (1705)...(1710)

&lt;223&gt; Polymorphism of 5 or 6 thymidine residues

&lt;221&gt; variation

&lt;222&gt; (487)...(487)

&lt;223&gt; Polymorphism of A or G

&lt;221&gt; variation

&lt;222&gt; (670)...(670)

&lt;223&gt; Polymorphism of C or T

&lt;400&gt; 3

```

gtggacttgt tgcagttgct gtaggattct aaatccaggt gattgtttca aactgagcat 60
caacaacaaa aacatttgta tgatatctat atttcaatc atg gac caa aat caa 114
                                   Met Asp Gln Asn Gln
                                   1 5

cat ttg aat aaa aca gca gag gca caa cct tca gag aat aag aaa aca 162
His Leu Asn Lys Thr Ala Glu Ala Gln Pro Ser Glu Asn Lys Lys Thr
                                   10 15 20

aga tac tgc aat gga ttg aag atg ttc ttg gca gct ctg tca ctc agc 210
Arg Tyr Cys Asn Gly Leu Lys Met Phe Leu Ala Ala Leu Ser Leu Ser
                                   25 30 35

ttt att gct aag aca cta ggt gca att att atg aaa agt tcc atc att 258
Phe Ile Ala Lys Thr Leu Gly Ala Ile Ile Met Lys Ser Ser Ile Ile
                                   40 45 50

cat ata gaa cgg aga ttt gag ata tcc tct tct ctt gtt ggt ttt att 306
His Ile Glu Arg Arg Phe Glu Ile Ser Ser Ser Leu Val Gly Phe Ile
                                   55 60 65

gac gga agc ttt gaa att gga aat ttg ctt gtg att gta ttt gtg agt 354
Asp Gly Ser Phe Glu Ile Gly Asn Leu Leu Val Ile Val Phe Val Ser
                                   70 75 80 85

tac ttt gga tcc aaa cta cat aga cca aag tta att gga atc ggt tgt 402
Tyr Phe Gly Ser Lys Leu His Arg Pro Lys Leu Ile Gly Ile Gly Cys
                                   90 95 100

```

ttc att atg gga att gga ggt gtt ttg act gct ttg cca cat ttc ttc 450  
 Phe Ile Met Gly Ile Gly Gly Val Leu Thr Ala Leu Pro His Phe Phe  
 105 110 115

atg gga tat tac agg tat tct aaa gaa act aat atc rat tca tca gaa 498  
 Met Gly Tyr Tyr Arg Tyr Ser Lys Glu Thr Asn Ile Xaa Ser Ser Glu  
 120 125 130

aat tca aca tcg acc tta tcc act tgt tta att aat caa att tta tca 546  
 Asn Ser Thr Ser Thr Leu Ser Thr Cys Leu Ile Asn Gln Ile Leu Ser  
 135 140 145

ctc aat aga gca tca cct gag ata gtg gga aaa ggt tgt tta aag gaa 594  
 Leu Asn Arg Ala Ser Pro Glu Ile Val Gly Lys Gly Cys Leu Lys Glu  
 150 155 160 165

tct ggg tca tac atg tgg ata tat gtg ttc atg ggt aat atg ctt cgt 642  
 Ser Gly Ser Tyr Met Trp Ile Tyr Val Phe Met Gly Asn Met Leu Arg  
 170 175 180

gga ata ggg gag act ccc ata gta cca ytg ggg ctt tct tac att gat 690  
 Gly Ile Gly Glu Thr Pro Ile Val Pro Xaa Gly Leu Ser Tyr Ile Asp  
 185 190 195

gat ttc gct aaa gaa gga cat tct tct ttg tat tta ggt ata ttg aat 738  
 Asp Phe Ala Lys Glu Gly His Ser Ser Leu Tyr Leu Gly Ile Leu Asn  
 200 205 210

gca ata gca atg att ggt cca atc att ggc ttt acc ctg gga tct ctg 786  
 Ala Ile Ala Met Ile Gly Pro Ile Ile Gly Phe Thr Leu Gly Ser Leu  
 215 220 225

ttt tct aaa atg tac gtg gat att gga tat gta gat cta agc act atc 834  
 Phe Ser Lys Met Tyr Val Asp Ile Gly Tyr Val Asp Leu Ser Thr Ile  
 230 235 240 245

agg ata act cct act gat tct cga tgg gtt gga gct tgg tgg ctt aat 882  
 Arg Ile Thr Pro Thr Asp Ser Arg Trp Val Gly Ala Trp Trp Leu Asn  
 250 255 260

ttc ctt gtg tct gga cta ttc tcc att att tct tcc ata cca ttc ttt 930  
 Phe Leu Val Ser Gly Leu Phe Ser Ile Ile Ser Ser Ile Pro Phe Phe  
 265 270 275

ttc ttg ccc caa act cca aat aaa cca caa aaa gaa aga aaa gct tca 978  
 Phe Leu Pro Gln Thr Pro Asn Lys Pro Gln Lys Glu Arg Lys Ala Ser  
 280 285 290

ctg tct ttg cat gtg ctg gaa aca aat gat gaa aag gat caa aca gct 1026  
 Leu Ser Leu His Val Leu Glu Thr Asn Asp Glu Lys Asp Gln Thr Ala  
 295 300 305

aat ttg acc aat caa gga aaa aat att acc aaa aat gtg act ggt ttt 1074  
 Asn Leu Thr Asn Gln Gly Lys Asn Ile Thr Lys Asn Val Thr Gly Phe  
 310 315 320 325

ttc cag tct ttt aaa agc atc ctt act aat ccc ctg tat gtt atg ttt 1122  
 Phe Gln Ser Phe Lys Ser Ile Leu Thr Asn Pro Leu Tyr Val Met Phe  
 330 335 340

gtg ctt ttg acg ttg tta caa gta agc agc tat att ggt gct ttt act 1170  
Val Leu Leu Thr Leu Leu Gln Val Ser Ser Tyr Ile Gly Ala Phe Thr  
345 350 355

tat gtc ttc aaa tac gta gag caa cag tat ggt cag cct tca tct aag 1218  
Tyr Val Phe Lys Tyr Val Glu Gln Tyr Gly Gln Pro Ser Ser Lys  
360 365 370

gct aac atc tta ttg gga gtc ata acc ata cct att ttt gca agt gga 1266  
Ala Asn Ile Leu Leu Gly Val Ile Thr Ile Pro Ile Phe Ala Ser Gly  
375 380 385

atg ttt tta gga gga tat atc att aaa aaa ttc aaa ctg aac acc gtt 1314  
Met Phe Leu Gly Gly Tyr Ile Ile Lys Lys Phe Lys Leu Asn Thr Val  
390 395 400 405

gga att gcc aaa ttc tca tgt ttt act gct gtg atg tca ttg tcc ttt 1362  
Gly Ile Ala Lys Phe Ser Cys Phe Thr Ala Val Met Ser Leu Ser Phe  
410 415 420

tac cta tta tat ttt ttc ata ctc tgt gaa aac aaa tca gtt gcc gga 1410  
Tyr Leu Leu Tyr Phe Phe Ile Leu Cys Glu Asn Lys Ser Val Ala Gly  
425 430 435

cta acc atg acc tat gat gga aat aat cca gtg aca tct cat aga gat 1458  
Leu Thr Met Thr Tyr Asp Gly Asn Asn Pro Val Thr Ser His Arg Asp  
440 445 450

gta cca ctt tct tat tgc aac tca gac tgc aat tgt gat gaa agt caa 1506  
Val Pro Leu Ser Tyr Cys Asn Ser Asp Cys Asn Cys Asp Glu Ser Gln  
455 460 465

tgg gaa cca gtc tgt gga aac aat gga ata act tac atc tca ccc tgt 1554  
Trp Glu Pro Val Cys Gly Asn Asn Gly Ile Thr Tyr Ile Ser Pro Cys  
470 475 480 485

cta gca ggt tgc aaa tct tca agt ggc aat aaa aag cct ata gtg ttt 1602  
Leu Ala Gly Cys Lys Ser Ser Ser Gly Asn Lys Lys Pro Ile Val Phe  
490 495 500

tac aac tgc agt tgt ttg gaa gta act ggt ctc cag aac aga aat tac 1650  
Tyr Asn Cys Ser Cys Leu Glu Val Thr Gly Leu Gln Asn Arg Asn Tyr  
505 510 515

tca gcc cat ttg ggt gaa tgc cca aga gat gat gct tgt aca agg aaa 1698  
Ser Ala His Leu Gly Glu Cys Pro Arg Asp Asp Ala Cys Thr Arg Lys  
520 525 530

ttt tac ttt ttg ttg caa tac aag tct tga a tttatttttc tctgcacttg 1749  
Phe Tyr Phe Leu Leu Gln Tyr Lys Ser \*  
535 540

gaggcacctc acatgtcatg ctgattgtta aaattgttca acctgaattg aaatcacttg 1809  
cactgggttt ccactcaatg gttatacgag cactaggagg aattctagct ccaatatatt 1869  
ttggggctct gattgataca acgtgtataa agtggtccac caacaactgt ggcacacgtg 1929  
ggcatgtag gacatataat tccacatcat tttcaagggt ctacttgggc ttgtcttcaa 1989  
tgttaagagt ctcactcactt gttttatata ttatattaat ttatgccatg aagaaaaaat 2049  
atcaagagaa agatatcaat gcatcagaaa atggaagtgt catggatgaa gcaaaccttag 2109  
aatccttaaa taataataaa cattttgtcc cttctgctgg ggcagatagt gaaacacatt 2169  
gttaagggga gaaaaaagc cacttctgct tctgtgtttc caaacagcat tgcattgatt 2229  
cagtaagatg ttatttttga ggagttcctg gtcctttcac taagaatttc cacatctttt 2289  
atggtggaag tataaataag cctatgaact tataataaaa caaactgtag gtagaaaaaa 2349

```

tgagagtact cattgttaca ttatagctac atatttgtgg ttaagggttag actatatgat 2409
ccatacaaat taaagtgaga gacatgggta ctgtgtaata aaagaaaaaa tacttgttca 2469
ggtaattcta attcttaata aaacaaatga gtatcataca ggtagagggtt aaaaaggagg 2529
agctagattc atattcctaag taaagagaaa tgcctagtgt ctattttatt aaacaaacaa 2589
acacagagtt tgaactataa tactaaggcc tgaagtctag cttggatata tgctacaata 2649
atatctgtta ctacacataaa attatatatt tcacagactt tatcaatgta taattaacaa 2709
ttatcttggt taagtaaatt tagaatacat ttaagtattg tggaagaaat aaagacattc 2769
caatatttgc aaaaaaaaaa aaaaaaaaaa 2799

```

<210> 4  
 <211> 542  
 <212> PRT  
 <213> H. sapiens

<220>  
 <221> VARIANT  
 <222> (130)...(130)  
 <223> Xaa = Asp or Asn

<221> VARIANT  
 <222> (191)...(191)  
 <223> Xaa = Leu

<400> 4

Met	Asp	Gln	Asn	Gln	His	Leu	Asn	Lys	Thr	Ala	Glu	Ala	Gln	Pro	Ser	1	5	10	15
Glu	Asn	Lys	Lys	Thr	Arg	Tyr	Cys	Asn	Gly	Leu	Lys	Met	Phe	Leu	Ala	20	25	30	
Ala	Leu	Ser	Leu	Ser	Phe	Ile	Ala	Lys	Thr	Leu	Gly	Ala	Ile	Ile	Met	35	40	45	
Lys	Ser	Ser	Ile	Ile	His	Ile	Glu	Arg	Arg	Phe	Glu	Ile	Ser	Ser	Ser	50	55	60	
Leu	Val	Gly	Phe	Ile	Asp	Gly	Ser	Phe	Glu	Ile	Gly	Asn	Leu	Leu	Val	65	70	75	80
Ile	Val	Phe	Val	Ser	Tyr	Phe	Gly	Ser	Lys	Leu	His	Arg	Pro	Lys	Leu	85	90	95	
Ile	Gly	Ile	Gly	Cys	Phe	Ile	Met	Gly	Ile	Gly	Gly	Val	Leu	Thr	Ala	100	105	110	
Leu	Pro	His	Phe	Phe	Met	Gly	Tyr	Tyr	Arg	Tyr	Ser	Lys	Glu	Thr	Asn	115	120	125	
Ile	Xaa	Ser	Ser	Glu	Asn	Ser	Thr	Ser	Thr	Leu	Ser	Thr	Cys	Leu	Ile	130	135	140	
Asn	Gln	Ile	Leu	Ser	Leu	Asn	Arg	Ala	Ser	Pro	Glu	Ile	Val	Gly	Lys	145	150	155	160
Gly	Cys	Leu	Lys	Glu	Ser	Gly	Ser	Tyr	Met	Trp	Ile	Tyr	Val	Phe	Met	165	170	175	
Gly	Asn	Met	Leu	Arg	Gly	Ile	Gly	Glu	Thr	Pro	Ile	Val	Pro	Xaa	Gly	180	185	190	
Leu	Ser	Tyr	Ile	Asp	Asp	Phe	Ala	Lys	Glu	Gly	His	Ser	Ser	Leu	Tyr	195	200	205	
Leu	Gly	Ile	Leu	Asn	Ala	Ile	Ala	Met	Ile	Gly	Pro	Ile	Ile	Gly	Phe	210	215	220	
Thr	Leu	Gly	Ser	Leu	Phe	Ser	Lys	Met	Tyr	Val	Asp	Ile	Gly	Tyr	Val	225	230	235	240
Asp	Leu	Ser	Thr	Ile	Arg	Ile	Thr	Pro	Thr	Asp	Ser	Arg	Trp	Val	Gly	245	250	255	
Ala	Trp	Trp	Leu	Asn	Phe	Leu	Val	Ser	Gly	Leu	Phe	Ser	Ile	Ile	Ser	260	265	270	
Ser	Ile	Pro	Phe	Phe	Phe	Leu	Pro	Gln	Thr	Pro	Asn	Lys	Pro	Gln	Lys	275	280	285	
Glu	Arg	Lys	Ala	Ser	Leu	Ser	Leu	His	Val	Leu	Glu	Thr	Asn	Asp	Glu	290	295	300	

Lys Asp Gln Thr Ala Asn Leu Thr Asn Gln Gly Lys Asn Ile Thr Lys  
 305 310 315 320  
 Asn Val Thr Gly Phe Phe Gln Ser Phe Lys Ser Ile Leu Thr Asn Pro  
 325 330 335  
 Leu Tyr Val Met Phe Val Leu Leu Thr Leu Gln Val Ser Ser Tyr  
 340 345 350  
 Ile Gly Ala Phe Thr Tyr Val Phe Lys Tyr Val Glu Gln Gln Tyr Gly  
 355 360 365  
 Gln Pro Ser Ser Lys Ala Asn Ile Leu Leu Gly Val Ile Thr Ile Pro  
 370 375 380  
 Ile Phe Ala Ser Gly Met Phe Leu Gly Gly Tyr Ile Ile Lys Lys Phe  
 385 390 395 400  
 Lys Leu Asn Thr Val Gly Ile Ala Lys Phe Ser Cys Phe Thr Ala Val  
 405 410 415  
 Met Ser Leu Ser Phe Tyr Leu Leu Tyr Phe Phe Ile Leu Cys Glu Asn  
 420 425 430  
 Lys Ser Val Ala Gly Leu Thr Met Thr Tyr Asp Gly Asn Asn Pro Val  
 435 440 445  
 Thr Ser His Arg Asp Val Pro Leu Ser Tyr Cys Asn Ser Asp Cys Asn  
 450 455 460  
 Cys Asp Glu Ser Gln Trp Glu Pro Val Cys Gly Asn Asn Gly Ile Thr  
 465 470 475 480  
 Tyr Ile Ser Pro Cys Leu Ala Gly Cys Lys Ser Ser Ser Gly Asn Lys  
 485 490 495  
 Lys Pro Ile Val Phe Tyr Asn Cys Ser Cys Leu Glu Val Thr Gly Leu  
 500 505 510  
 Gln Asn Arg Asn Tyr Ser Ala His Leu Gly Glu Cys Pro Arg Asp Asp  
 515 520 525  
 Ala Cys Thr Arg Lys Phe Tyr Phe Leu Leu Gln Tyr Lys Ser  
 530 535 540

&lt;210&gt; 5

&lt;211&gt; 2800

&lt;212&gt; DNA

&lt;213&gt; H. sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (100)...(2175)

&lt;223&gt; Coding sequence of ATnov3.1

&lt;221&gt; variation

&lt;222&gt; (1705)...(1710)

&lt;223&gt; Polymorphism of 5 or 6 T residues.

&lt;221&gt; variation

&lt;222&gt; (487)...(487)

&lt;223&gt; Polymorphism of A or G

&lt;221&gt; variation

&lt;222&gt; (670)...(670)

&lt;223&gt; Polymorphism of C or T

&lt;400&gt; 5

gtggacttggt tgcagttgct gtaggattct aaatccaggt gattgtttca aactgagcat 60  
 caacaacaaa aacatttgta tgatatctat atttcaatc atg gac caa aat caa 114  
 Met Asp Gln Asn Gln  
 1 5

cat ttg aat aaa aca gca gag gca caa cct tca gag aat aag aaa aca 162  
 His Leu Asn Lys Thr Ala Glu Ala Gln Pro Ser Glu Asn Lys Lys Thr  
 10 15 20

aga tac tgc aat gga ttg aag atg ttc ttg gca gct ctg tca ctc agc Arg Tyr Cys Asn Gly Leu Lys Met Phe Leu Ala Ala Leu Ser Leu Ser 25 30 35	210
ttt att gct aag aca cta ggt gca att att atg aaa agt tcc atc att Phe Ile Ala Lys Thr Leu Gly Ala Ile Ile Met Lys Ser Ser Ile Ile 40 45 50	258
cat ata gaa cgg aga ttt gag ata tcc tct tct ctt gtt ggt ttt att His Ile Glu Arg Arg Phe Glu Ile Ser Ser Ser Leu Val Gly Phe Ile 55 60 65	306
gac gga agc ttt gaa att gga aat ttg ctt gtg att gta ttt gtg agt Asp Gly Ser Phe Glu Ile Gly Asn Leu Leu Val Ile Val Phe Val Ser 70 75 80 85	354
tac ttt gga tcc aaa cta cat aga cca aag tta att gga atc ggt tgt Tyr Phe Gly Ser Lys Leu His Arg Pro Lys Leu Ile Gly Ile Gly Cys 90 95 100	402
ttc att atg gga att gga ggt gtt ttg act gct ttg cca cat ttc ttc Phe Ile Met Gly Ile Gly Gly Val Leu Thr Ala Leu Pro His Phe Phe 105 110 115	450
atg gga tat tac agg tat tct aaa gaa act aat atc rat tca tca gaa Met Gly Tyr Tyr Arg Tyr Ser Lys Glu Thr Asn Ile Xaa Ser Ser Glu 120 125 130	498
aat tca aca tgc acc tta tcc act tgt tta att aat caa att tta tca Asn Ser Thr Ser Thr Leu Ser Thr Cys Leu Ile Asn Gln Ile Leu Ser 135 140 145	546
ctc aat aga gca tca cct gag ata gtg gga aaa ggt tgt tta aag gaa Leu Asn Arg Ala Ser Pro Glu Ile Val Gly Lys Gly Cys Leu Lys Glu 150 155 160 165	594
tct ggg tca tac atg tgg ata tat gtg ttc atg ggt aat atg ctt cgt Ser Gly Ser Tyr Met Trp Ile Tyr Val Phe Met Gly Asn Met Leu Arg 170 175 180	642
gga ata ggg gag act ccc ata gta cca ytg ggg ctt tct tac att gat Gly Ile Gly Glu Thr Pro Ile Val Pro Xaa Gly Leu Ser Tyr Ile Asp 185 190 195	690
gat ttc gct aaa gaa gga cat tct tct ttg tat tta ggt ata ttg aat Asp Phe Ala Lys Glu Gly His Ser Ser Leu Tyr Leu Gly Ile Leu Asn 200 205 210	738
gca ata gca atg att ggt cca atc att ggc ttt acc ctg gga tct ctg Ala Ile Ala Met Ile Gly Pro Ile Ile Gly Phe Thr Leu Gly Ser Leu 215 220 225	786
ttt tct aaa atg tac gtg gat att gga tat gta gat cta agc act atc Phe Ser Lys Met Tyr Val Asp Ile Gly Tyr Val Asp Leu Ser Thr Ile 230 235 240 245	834
agg ata act cct act gat tct cga tgg gtt gga gct tgg tgg ctt aat Arg Ile Thr Pro Thr Asp Ser Arg Trp Val Gly Ala Trp Trp Leu Asn 250 255 260	882

ttc ctt gtg tct gga cta ttc tcc att att tct tcc ata cca ttc ttt Phe Leu Val Ser Gly Leu Phe Ser Ile Ile Ser Ser Ile Pro Phe Phe 265 270 275	930
ttc ttg ccc caa act cca aat aaa cca caa aaa gaa aga aaa gct tca Phe Leu Pro Gln Thr Pro Asn Lys Pro Gln Lys Glu Arg Lys Ala Ser 280 285 290	978
ctg tct ttg cat gtg ctg gaa aca aat gat gaa aag gat caa aca gct Leu Ser Leu His Val Leu Glu Thr Asn Asp Glu Lys Asp Gln Thr Ala 295 300 305	1026
aat ttg acc aat caa gga aaa aat att acc aaa aat gtg act ggt ttt Asn Leu Thr Asn Gln Gly Lys Asn Ile Thr Lys Asn Val Thr Gly Phe 310 315 320 325	1074
ttc cag tct ttt aaa agc atc ctt act aat ccc ctg tat gtt atg ttt Phe Gln Ser Phe Lys Ser Ile Leu Thr Asn Pro Leu Tyr Val Met Phe 330 335 340	1122
gtg ctt ttg acg ttg tta caa gta agc agc tat att ggt gct ttt act Val Leu Leu Thr Leu Leu Gln Val Ser Ser Tyr Ile Gly Ala Phe Thr 345 350 355	1170
tat gtc ttc aaa tac gta gag caa cag tat ggt cag cct tca tct aag Tyr Val Phe Lys Tyr Val Glu Gln Gln Tyr Gly Gln Pro Ser Ser Lys 360 365 370	1218
gct aac atc tta ttg gga gtc ata acc ata cct att ttt gca agt gga Ala Asn Ile Leu Leu Gly Val Ile Thr Ile Pro Ile Phe Ala Ser Gly 375 380 385	1266
atg ttt tta gga gga tat atc att aaa aaa ttc aaa ctg aac acc gtt Met Phe Leu Gly Gly Tyr Ile Ile Lys Lys Phe Lys Leu Asn Thr Val 390 395 400 405	1314
gga att gcc aaa ttc tca tgt ttt act gct gtg atg tca ttg tcc ttt Gly Ile Ala Lys Phe Ser Cys Phe Thr Ala Val Met Ser Leu Ser Phe 410 415 420	1362
tac cta tta tat ttt ttc ata ctc tgt gaa aac aaa tca gtt gcc gga Tyr Leu Leu Tyr Phe Phe Ile Leu Cys Glu Asn Lys Ser Val Ala Gly 425 430 435	1410
cta acc atg acc tat gat gga aat aat cca gtg aca tct cat aga gat Leu Thr Met Thr Tyr Asp Gly Asn Asn Pro Val Thr Ser His Arg Asp 440 445 450	1458
gta cca ctt tct tat tgc aac tca gac tgc aat tgt gat gaa agt caa Val Pro Leu Ser Tyr Cys Asn Ser Asp Cys Asn Cys Asp Glu Ser Gln 455 460 465	1506
tggt gaa cca gtc tgt gga aac aat gga ata act tac atc tca ccc tgt Trp Glu Pro Val Cys Gly Asn Asn Gly Ile Thr Tyr Ile Ser Pro Cys 470 475 480 485	1554
cta gca ggt tgc aaa tct tca agt ggc aat aaa aag cct ata gtg ttt Leu Ala Gly Cys Lys Ser Ser Ser Gly Asn Lys Lys Pro Ile Val Phe 490 495 500	1602



tac aac tgc agt tgt ttg gaa gta act ggt ctc cag aac aga aat tac 1650  
 Tyr Asn Cys Ser Cys Leu Glu Val Thr Gly Leu Gln Asn Arg Asn Tyr  
 505 510 515

tca gcc cat ttg ggt gaa tgc cca aga gat gat gct tgt aca agg aaa 1698  
 Ser Ala His Leu Gly Glu Cys Pro Arg Asp Asp Ala Cys Thr Arg Lys  
 520 525 530

ttt tac ttt ttt gtt gca ata caa gtc ttg aat tta ttt ttc tct gca 1746  
 Phe Tyr Phe Phe Val Ala Ile Gln Val Leu Asn Leu Phe Phe Ser Ala  
 535 540 545

ctt gga ggc acc tca cat gtc atg ctg att gtt aaa att gtt caa cct 1794  
 Leu Gly Gly Thr Ser His Val Met Leu Ile Val Lys Ile Val Gln Pro  
 550 555 560 565

gaa ttg aaa tca ctt gca ctg ggt ttc cac tca atg gtt ata cga gca 1842  
 Glu Leu Lys Ser Leu Ala Leu Gly Phe His Ser Met Val Ile Arg Ala  
 570 575 580

cta gga gga att cta gct cca ata tat ttt ggg gct ctg att gat aca 1890  
 Leu Gly Gly Ile Leu Ala Pro Ile Tyr Phe Gly Ala Leu Ile Asp Thr  
 585 590 595

acg tgt ata aag tgg tcc acc aac aac tgt ggc aca cgt ggg tca tgt 1938  
 Thr Cys Ile Lys Trp Ser Thr Asn Asn Cys Gly Thr Arg Gly Ser Cys  
 600 605 610

agg aca tat aat tcc aca tca ttt tca agg gtc tac ttg ggc ttg tct 1986  
 Arg Thr Tyr Asn Ser Thr Ser Phe Ser Arg Val Tyr Leu Gly Leu Ser  
 615 620 625

tca atg tta aga gtc tca tca ctt gtt tta tat att ata tta att tat 2034  
 Ser Met Leu Arg Val Ser Ser Leu Val Leu Tyr Ile Ile Leu Ile Tyr  
 630 635 640 645

gcc atg aag aaa aaa tat caa gag aaa gat atc aat gca tca gaa aat 2082  
 Ala Met Lys Lys Lys Tyr Gln Glu Lys Asp Ile Asn Ala Ser Glu Asn  
 650 655 660

gga agt gtc atg gat gaa gca aac tta gaa tcc tta aat aaa aat aaa 2130  
 Gly Ser Val Met Asp Glu Ala Asn Leu Glu Ser Leu Asn Lys Asn Lys  
 665 670 675

cat ttt gtc cct tct gct ggg gca gat agt gaa aca cat tgt taa 2175  
 His Phe Val Pro Ser Ala Gly Ala Asp Ser Glu Thr His Cys \*  
 680 685 690

ggggagaaaa aaagccactt ctgcttctgt gtttccaaac agcattgcat tgattcagta 2235  
 agatgttatt ttgaggagt tcctggctct ttactaaga atttccacat cttttatggt 2295  
 ggaagtataa ataagcctat gaacttataa taaaacaaac tgtaggtaga aaaaatgaga 2355  
 gtactcattg ttacattata gctacatatt tgtgggtaag gttagactat atgatccata 2415  
 caaattaaag tgagagacat ggttactgtg taataaaaaga aaaaataactt gttcaggtaa 2475  
 ttctaattct taataaaaaca aatgagtatc atacaggtag aggttaaaaaa ggaggagcta 2535  
 gattcatatc ctaagtaaag agaaatgcct agtgtctatt ttattaaaca aacaaacaca 2595  
 gagtttgaac tataatacta aggcctgaag tctagcttgg atatattgcta caataatc 2655  
 tggtactcac ataaaattat atatttcaca gactttatca atgtataatt aacaattatc 2715  
 ttgtttaagt aaatttagaa tacatttaag tattgtggaa gaaataaaga cattccaata 2775  
 tttgcaaaaaa aaaaaaaaaa aaaaa 2800

&lt;210&gt; 6

&lt;211&gt; 691

<212> PRT  
 <213> H. sapiens

<220>  
 <221> VARIANT  
 <222> (130)...(130)  
 <223> Xaa = Asp or Asn

<221> VARIANT  
 <222> (191)...(191)  
 <223> Xaa = Leu

<400> 6  
 Met Asp Gln Asn Gln His Leu Asn Lys Thr Ala Glu Ala Gln Pro Ser  
 1 5 10 15  
 Glu Asn Lys Lys Thr Arg Tyr Cys Asn Gly Leu Lys Met Phe Leu Ala  
 20 25 30  
 Ala Leu Ser Leu Ser Phe Ile Ala Lys Thr Leu Gly Ala Ile Ile Met  
 35 40 45  
 Lys Ser Ser Ile Ile His Ile Glu Arg Arg Phe Glu Ile Ser Ser Ser  
 50 55 60  
 Leu Val Gly Phe Ile Asp Gly Ser Phe Glu Ile Gly Asn Leu Leu Val  
 65 70 75 80  
 Ile Val Phe Val Ser Tyr Phe Gly Ser Lys Leu His Arg Pro Lys Leu  
 85 90 95  
 Ile Gly Ile Gly Cys Phe Ile Met Gly Ile Gly Gly Val Leu Thr Ala  
 100 105 110  
 Leu Pro His Phe Phe Met Gly Tyr Tyr Arg Tyr Ser Lys Glu Thr Asn  
 115 120 125  
 Ile Xaa Ser Ser Glu Asn Ser Thr Ser Thr Leu Ser Thr Cys Leu Ile  
 130 135 140  
 Asn Gln Ile Leu Ser Leu Asn Arg Ala Ser Pro Glu Ile Val Gly Lys  
 145 150 155 160  
 Gly Cys Leu Lys Glu Ser Gly Ser Tyr Met Trp Ile Tyr Val Phe Met  
 165 170 175  
 Gly Asn Met Leu Arg Gly Ile Gly Glu Thr Pro Ile Val Pro Xaa Gly  
 180 185 190  
 Leu Ser Tyr Ile Asp Asp Phe Ala Lys Glu Gly His Ser Ser Leu Tyr  
 195 200 205  
 Leu Gly Ile Leu Asn Ala Ile Ala Met Ile Gly Pro Ile Ile Gly Phe  
 210 215 220  
 Thr Leu Gly Ser Leu Phe Ser Lys Met Tyr Val Asp Ile Gly Tyr Val  
 225 230 235 240  
 Asp Leu Ser Thr Ile Arg Ile Thr Pro Thr Asp Ser Arg Trp Val Gly  
 245 250 255  
 Ala Trp Trp Leu Asn Phe Leu Val Ser Gly Leu Phe Ser Ile Ile Ser  
 260 265 270  
 Ser Ile Pro Phe Phe Phe Leu Pro Gln Thr Pro Asn Lys Pro Gln Lys  
 275 280 285  
 Glu Arg Lys Ala Ser Leu Ser Leu His Val Leu Glu Thr Asn Asp Glu  
 290 295 300  
 Lys Asp Gln Thr Ala Asn Leu Thr Asn Gln Gly Lys Asn Ile Thr Lys  
 305 310 315 320  
 Asn Val Thr Gly Phe Phe Gln Ser Phe Lys Ser Ile Leu Thr Asn Pro  
 325 330 335  
 Leu Tyr Val Met Phe Val Leu Leu Thr Leu Leu Gln Val Ser Ser Tyr  
 340 345 350  
 Ile Gly Ala Phe Thr Tyr Val Phe Lys Tyr Val Glu Gln Gln Tyr Gly  
 355 360 365  
 Gln Pro Ser Ser Lys Ala Asn Ile Leu Leu Gly Val Ile Thr Ile Pro  
 370 375 380

```

Ile Phe Ala Ser Gly Met Phe Leu Gly Gly Tyr Ile Ile Lys Lys Phe
385                      390                      395                      400
Lys Leu Asn Thr Val Gly Ile Ala Lys Phe Ser Cys Phe Thr Ala Val
                      405                      410                      415
Met Ser Leu Ser Phe Tyr Leu Leu Tyr Phe Phe Ile Leu Cys Glu Asn
                      420                      425                      430
Lys Ser Val Ala Gly Leu Thr Met Thr Tyr Asp Gly Asn Asn Pro Val
                      435                      440                      445
Thr Ser His Arg Asp Val Pro Leu Ser Tyr Cys Asn Ser Asp Cys Asn
                      450                      455                      460
Cys Asp Glu Ser Gln Trp Glu Pro Val Cys Gly Asn Asn Gly Ile Thr
465                      470                      475                      480
Tyr Ile Ser Pro Cys Leu Ala Gly Cys Lys Ser Ser Ser Gly Asn Lys
                      485                      490                      495
Lys Pro Ile Val Phe Tyr Asn Cys Ser Cys Leu Glu Val Thr Gly Leu
                      500                      505                      510
Gln Asn Arg Asn Tyr Ser Ala His Leu Gly Glu Cys Pro Arg Asp Asp
                      515                      520                      525
Ala Cys Thr Arg Lys Phe Tyr Phe Phe Val Ala Ile Gln Val Leu Asn
530                      535                      540
Leu Phe Phe Ser Ala Leu Gly Gly Thr Ser His Val Met Leu Ile Val
545                      550                      555                      560
Lys Ile Val Gln Pro Glu Leu Lys Ser Leu Ala Leu Gly Phe His Ser
                      565                      570                      575
Met Val Ile Arg Ala Leu Gly Gly Ile Leu Ala Pro Ile Tyr Phe Gly
                      580                      585                      590
Ala Leu Ile Asp Thr Thr Cys Ile Lys Trp Ser Thr Asn Asn Cys Gly
                      595                      600                      605
Thr Arg Gly Ser Cys Arg Thr Tyr Asn Ser Thr Ser Phe Ser Arg Val
610                      615                      620
Tyr Leu Gly Leu Ser Ser Met Leu Arg Val Ser Ser Leu Val Leu Tyr
625                      630                      635                      640
Ile Ile Leu Ile Tyr Ala Met Lys Lys Lys Tyr Gln Glu Lys Asp Ile
                      645                      650                      655
Asn Ala Ser Glu Asn Gly Ser Val Met Asp Glu Ala Asn Leu Glu Ser
660                      665                      670
Leu Asn Lys Asn Lys His Phe Val Pro Ser Ala Gly Ala Asp Ser Glu
675                      680                      685
Thr His Cys
690

```

&lt;210&gt; 7

&lt;211&gt; 2360

&lt;212&gt; DNA

&lt;213&gt; H. sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (100)...(1729)

&lt;223&gt; Coding sequence ATnov3.2

&lt;221&gt; variation

&lt;222&gt; (1705)...(1710)

&lt;223&gt; Polymorphism of 5 or 6 T residues

&lt;221&gt; variation

&lt;222&gt; (487)...(487)

&lt;223&gt; Polymorphism of A or G

&lt;221&gt; variation

&lt;222&gt; (670)...(670)

&lt;223&gt; Polymorphism of C or T

<400> 7  
gtggacttgt tgcagttgct gtaggattct aaatccagggt gattgtttca aactgagcat 60  
caacaacaaa aacatttgta tgatatctat atttcaatc atg gac caa aat caa 114  
Met Asp Gln Asn Gln  
1 5

cat ttg aat aaa aca gca gag gca caa cct tca gag aat aag aaa aca 162  
His Leu Asn Lys Thr Ala Glu Ala Gln Pro Ser Glu Asn Lys Lys Thr  
10 15 20

aga tac tgc aat gga ttg aag atg ttc ttg gca gct ctg tca ctc agc 210  
Arg Tyr Cys Asn Gly Leu Lys Met Phe Leu Ala Ala Leu Ser Leu Ser  
25 30 35

ttt att gct aag aca cta ggt gca att att atg aaa agt tcc atc att 258  
Phe Ile Ala Lys Thr Leu Gly Ala Ile Ile Met Lys Ser Ser Ile Ile  
40 45 50

cat ata gaa cgg aga ttt gag ata tcc tct tct ctt gtt ggt ttt att 306  
His Ile Glu Arg Arg Phe Glu Ile Ser Ser Ser Leu Val Gly Phe Ile  
55 60 65

gac gga agc ttt gaa att gga aat ttg ctt gtg att gta ttt gtg agt 354  
Asp Gly Ser Phe Glu Ile Gly Asn Leu Leu Val Ile Val Phe Val Ser  
70 75 80 85

tac ttt gga tcc aaa cta cat aga cca aag tta att gga atc ggt tgt 402  
Tyr Phe Gly Ser Lys Leu His Arg Pro Lys Leu Ile Gly Ile Gly Cys  
90 95 100

ttc att atg gga att gga ggt gtt ttg act gct ttg cca cat ttc ttc 450  
Phe Ile Met Gly Ile Gly Gly Val Leu Thr Ala Leu Pro His Phe Phe  
105 110 115

atg gga tat tac agg tat tct aaa gaa act aat atc rat tca tca gaa 498  
Met Gly Tyr Tyr Arg Tyr Ser Lys Glu Thr Asn Ile Xaa Ser Ser Glu  
120 125 130

aat tca aca tcg acc tta tcc act tgt tta att aat caa att tta tca 546  
Asn Ser Thr Ser Thr Leu Ser Thr Cys Leu Ile Asn Gln Ile Leu Ser  
135 140 145

ctc aat aga gca tca cct gag ata gtg gga aaa ggt tgt tta aag gaa 594  
Leu Asn Arg Ala Ser Pro Glu Ile Val Gly Lys Gly Cys Leu Lys Glu  
150 155 160 165

tct ggg tca tac atg tgg ata tat gtg ttc atg ggt aat atg ctt cgt 642  
Ser Gly Ser Tyr Met Trp Ile Tyr Val Phe Met Gly Asn Met Leu Arg  
170 175 180

gga ata ggg gag act ccc ata gta cca ytg ggg ctt tct tac att gat 690  
Gly Ile Gly Glu Thr Pro Ile Val Pro Xaa Gly Leu Ser Tyr Ile Asp  
185 190 195

gat ttc gct aaa gaa gga cat tct tct ttg tat tta ggt ata ttg aat 738  
Asp Phe Ala Lys Glu Gly His Ser Ser Leu Tyr Leu Gly Ile Leu Asn  
200 205 210

gca ata gca atg att ggt cca atc att ggc ttt acc ctg gga tct ctg 786  
Ala Ile Ala Met Ile Gly Pro Ile Ile Gly Phe Thr Leu Gly Ser Leu  
215 220 225

ttt tct aaa atg tac gtg gat att gga tat gta gat cta agc act atc Phe Ser Lys Met Tyr Val Asp Ile Gly Tyr Val Asp Leu Ser Thr Ile 230 235 240 245	834
agg ata act cct act gat tct cga tgg gtt gga gct tgg tgg ctt aat Arg Ile Thr Pro Thr Asp Ser Arg Trp Val Gly Ala Trp Trp Leu Asn 250 255 260	882
ttc ctt gtg tct gga cta ttc tcc att att tct tcc ata cca ttc ttt Phe Leu Val Ser Gly Leu Phe Ser Ile Ile Ser Ser Ile Pro Phe Phe 265 270 275	930
ttc ttg ccc caa act cca aat aaa cca caa aaa gaa aga aaa gct tca Phe Leu Pro Gln Thr Pro Asn Lys Pro Gln Lys Glu Arg Lys Ala Ser 280 285 290	978
ctg tct ttg cat gtg ctg gaa aca aat gat gaa aag gat caa aca gct Leu Ser Leu His Val Leu Glu Thr Asn Asp Glu Lys Asp Gln Thr Ala 295 300 305	1026
aat ttg acc aat caa gga aaa aat att acc aaa aat gtg act ggt ttt Asn Leu Thr Asn Gln Gly Lys Asn Ile Thr Lys Asn Val Thr Gly Phe 310 315 320 325	1074
ttc cag tct ttt aaa agc atc ctt act aat ccc ctg tat gtt atg ttt Phe Gln Ser Phe Lys Ser Ile Leu Thr Asn Pro Leu Tyr Val Met Phe 330 335 340	1122
gtg ctt ttg acg ttg tta caa gta agc agc tat att ggt gct ttt act Val Leu Leu Thr Leu Leu Gln Val Ser Ser Tyr Ile Gly Ala Phe Thr 345 350 355	1170
tat gtc ttc aaa tac gta gag caa cag tat ggt cag cct tca tct aag Tyr Val Phe Lys Tyr Val Glu Gln Gln Tyr Gly Gln Pro Ser Ser Lys 360 365 370	1218
gct aac atc tta ttg gga gtc ata acc ata cct att ttt gca agt gga Ala Asn Ile Leu Leu Gly Val Ile Thr Ile Pro Ile Phe Ala Ser Gly 375 380 385	1266
atg ttt tta gga gga tat atc att aaa aaa ttc aaa ctg aac acc gtt Met Phe Leu Gly Gly Tyr Ile Ile Lys Lys Phe Lys Leu Asn Thr Val 390 395 400 405	1314
gga att gcc aaa ttc tca tgt ttt act gct gtg atg tca ttg tcc ttt Gly Ile Ala Lys Phe Ser Cys Phe Thr Ala Val Met Ser Leu Ser Phe 410 415 420	1362
tac cta tta tat ttt ttc ata ctc tgt gaa aac aaa tca gtt gcc gga Tyr Leu Leu Tyr Phe Phe Ile Leu Cys Glu Asn Lys Ser Val Ala Gly 425 430 435	1410
cta acc atg acc tat gat gga aat aat cca gtg aca tct cat aga gat Leu Thr Met Thr Tyr Asp Gly Asn Asn Pro Val Thr Ser His Arg Asp 440 445 450	1458
gta cca ctt tct tat tgc aac tca gac tgc aat tgt gat gaa agt caa Val Pro Leu Ser Tyr Cys Asn Ser Asp Cys Asn Cys Asp Glu Ser Gln 455 460 465	1506

```

tgg gaa cca gtc tgt gga aac aat gga ata act tac atc tca ccc tgt      1554
Trp Glu Pro Val Cys Gly Asn Asn Gly Ile Thr Tyr Ile Ser Pro Cys
470                               475                               480                               485

cta gca ggt tgc aaa tct tca agt ggc aat aaa aag cct ata gtg ttt      1602
Leu Ala Gly Cys Lys Ser Ser Ser Gly Asn Lys Lys Pro Ile Val Phe
                               490                               495                               500

tac aac tgc agt tgt ttg gaa gta act ggt ctc cag aac aga aat tac      1650
Tyr Asn Cys Ser Cys Leu Glu Val Thr Gly Leu Gln Asn Arg Asn Tyr
                               505                               510                               515

tca gcc cat ttg ggt gaa tgc cca aga gat gat gct tgt aca agg aaa      1698
Ser Ala His Leu Gly Glu Cys Pro Arg Asp Asp Ala Cys Thr Arg Lys
                               520                               525                               530

ttt tac ttt ttg ttg caa tac aag tct tga a tttatttttc tctgcacttg      1749
Phe Tyr Phe Leu Leu Gln Tyr Lys Ser *
                               535                               540

gaggcacctc acatgtcatg ctgattgtta aaattgttca acctgaattg aaatcacttg      1809
cactgggttt ccactcaatg gttatacgag cactaggagg aattctagct ccaatatatt      1869
ttggggctct gattgataca acgtgtataa agtgggtccac caacaactgt ggcacacgtg      1929
ggatcatgtag gacatataat tccacatcat ttccaagggt ctacttgggc ttgtcttcaa      1989
tgtaaagagt ctcactcactt gttttatata ttatattaat ttatgccatg aagaaaaaat      2049
atcaagagaa agatatcaat gcatcagaaa atggaagtgt catggatgaa gcaaacttag      2109
aatccttaaa taaaaataaa cattttgtcc cttctgctgg ggcagatagt gaaacacatt      2169
gttaagggga gaaaaaaaag cacttctgct tctgtgtttc caaacagcat tgcattgatt      2229
cagtaagatg ttatttttga ggagttcctg gtcctttcac taagaatttc cacatctttt      2289
atggtggaag tataataaag cctatgaact tataataaaa caaactgtag gtagaaaaaa      2349
aaaaaaaaa a                               2360

```

&lt;210&gt; 8

&lt;211&gt; 542

&lt;212&gt; PRT

&lt;213&gt; H. sapiens

&lt;220&gt;

&lt;221&gt; VARIANT

&lt;222&gt; (130)...(130)

&lt;223&gt; Xaa = Asp or Asn

&lt;221&gt; VARIANT

&lt;222&gt; (191)...(191)

&lt;223&gt; Xaa = Leu

&lt;400&gt; 8

```

Met Asp Gln Asn Gln His Leu Asn Lys Thr Ala Glu Ala Gln Pro Ser
 1           5           10           15
Glu Asn Lys Lys Thr Arg Tyr Cys Asn Gly Leu Lys Met Phe Leu Ala
          20          25          30
Ala Leu Ser Leu Ser Phe Ile Ala Lys Thr Leu Gly Ala Ile Ile Met
          35          40          45
Lys Ser Ser Ile Ile His Ile Glu Arg Arg Phe Glu Ile Ser Ser Ser
          50          55          60
Leu Val Gly Phe Ile Asp Gly Ser Phe Glu Ile Gly Asn Leu Leu Val
          65          70          75          80
Ile Val Phe Val Ser Tyr Phe Gly Ser Lys Leu His Arg Pro Lys Leu
          85          90          95
Ile Gly Ile Gly Cys Phe Ile Met Gly Ile Gly Gly Val Leu Thr Ala
          100          105          110

```

Leu Pro His Phe Phe Met Gly Tyr Tyr Arg Tyr Ser Lys Glu Thr Asn  
 115 120 125  
 Ile Xaa Ser Ser Glu Asn Ser Thr Ser Thr Leu Ser Thr Cys Leu Ile  
 130 135 140  
  
 Asn Gln Ile Leu Ser Leu Asn Arg Ala Ser Pro Glu Ile Val Gly Lys  
 145 150 155 160  
 Gly Cys Leu Lys Glu Ser Gly Ser Tyr Met Trp Ile Tyr Val Phe Met  
 165 170 175  
 Gly Asn Met Leu Arg Gly Ile Gly Glu Thr Pro Ile Val Pro Xaa Gly  
 180 185 190  
 Leu Ser Tyr Ile Asp Asp Phe Ala Lys Glu Gly His Ser Ser Leu Tyr  
 195 200 205  
 Leu Gly Ile Leu Asn Ala Ile Ala Met Ile Gly Pro Ile Ile Gly Phe  
 210 215 220  
 Thr Leu Gly Ser Leu Phe Ser Lys Met Tyr Val Asp Ile Gly Tyr Val  
 225 230 235 240  
 Asp Leu Ser Thr Ile Arg Ile Thr Pro Thr Asp Ser Arg Trp Val Gly  
 245 250 255  
 Ala Trp Trp Leu Asn Phe Leu Val Ser Gly Leu Phe Ser Ile Ile Ser  
 260 265 270  
 Ser Ile Pro Phe Phe Leu Pro Gln Thr Pro Asn Lys Pro Gln Lys  
 275 280 285  
 Glu Arg Lys Ala Ser Leu Ser Leu His Val Leu Glu Thr Asn Asp Glu  
 290 295 300  
 Lys Asp Gln Thr Ala Asn Leu Thr Asn Gln Gly Lys Asn Ile Thr Lys  
 305 310 315 320  
 Asn Val Thr Gly Phe Phe Gln Ser Phe Lys Ser Ile Leu Thr Asn Pro  
 325 330 335  
 Leu Tyr Val Met Phe Val Leu Leu Thr Leu Leu Gln Val Ser Ser Tyr  
 340 345 350  
 Ile Gly Ala Phe Thr Tyr Val Phe Lys Tyr Val Glu Gln Gln Tyr Gly  
 355 360 365  
 Gln Pro Ser Ser Lys Ala Asn Ile Leu Leu Gly Val Ile Thr Ile Pro  
 370 375 380  
 Ile Phe Ala Ser Gly Met Phe Leu Gly Gly Tyr Ile Ile Lys Lys Phe  
 385 390 395 400  
 Lys Leu Asn Thr Val Gly Ile Ala Lys Phe Ser Cys Phe Thr Ala Val  
 405 410 415  
 Met Ser Leu Ser Phe Tyr Leu Leu Tyr Phe Phe Ile Leu Cys Glu Asn  
 420 425 430  
 Lys Ser Val Ala Gly Leu Thr Met Thr Tyr Asp Gly Asn Asn Pro Val  
 435 440 445  
 Thr Ser His Arg Asp Val Pro Leu Ser Tyr Cys Asn Ser Asp Cys Asn  
 450 455 460  
 Cys Asp Glu Ser Gln Trp Glu Pro Val Cys Gly Asn Asn Gly Ile Thr  
 465 470 475 480  
 Tyr Ile Ser Pro Cys Leu Ala Gly Cys Lys Ser Ser Ser Gly Asn Lys  
 485 490 495  
 Lys Pro Ile Val Phe Tyr Asn Cys Ser Cys Leu Glu Val Thr Gly Leu  
 500 505 510  
 Gln Asn Arg Asn Tyr Ser Ala His Leu Gly Glu Cys Pro Arg Asp Asp  
 515 520 525  
 Ala Cys Thr Arg Lys Phe Tyr Phe Leu Leu Gln Tyr Lys Ser  
 530 535 540

<210> 9  
 <211> 2361  
 <212> DNA  
 <213> H. sapiens

<220>

<400> 9															60
gtggacttgt tgcagttgct gtaggattct aaatccaggt gattgtttca aactgagcat															114
caacaacaaa aacatttgta tgatatctat atttcaatc atg gac caa aat caa															
Met Asp Gln Asn Gln															
1 5															
cat ttg aat aaa aca gca gag gca caa cct tca gag aat aag aaa aca															162
His Leu Asn Lys Thr Ala Glu Ala Gln Pro Ser Glu Asn Lys Lys Thr															
10 15 20															
aga tac tgc aat gga ttg aag atg ttc ttg gca gct ctg tca ctc agc															210
Arg Tyr Cys Asn Gly Leu Lys Met Phe Leu Ala Ala Leu Ser Leu Ser															
25 30 35															
ttt att gct aag aca cta ggt gca att att atg aaa agt tcc atc att															258
Phe Ile Ala Lys Thr Leu Gly Ala Ile Ile Met Lys Ser Ser Ile Ile															
40 45 50															
cat ata gaa cgg aga ttt gag ata tcc tct tct ctt gtt ggt ttt att															306
His Ile Glu Arg Arg Phe Glu Ile Ser Ser Ser Leu Val Gly Phe Ile															
55 60 65															
gac gga agc ttt gaa att gga aat ttg ctt gtg att gta ttt gtg agt															354
Asp Gly Ser Phe Glu Ile Gly Asn Leu Leu Val Ile Val Phe Val Ser															
70 75 80 85															
tac ttt gga tcc aaa cta cat aga cca aag tta att gga atc ggt tgt															402
Tyr Phe Gly Ser Lys Leu His Arg Pro Lys Leu Ile Gly Ile Gly Cys															
90 95 100															
ttc att atg gga att gga ggt gtt ttg act gct ttg cca cat ttc ttc															450
Phe Ile Met Gly Ile Gly Gly Val Leu Thr Ala Leu Pro His Phe Phe															
105 110 115															
atg gga tat tac agg tat tct aaa gaa act aat atc rat tca tca gaa															498
Met Gly Tyr Tyr Arg Tyr Ser Lys Glu Thr Asn Ile Xaa Ser Ser Glu															
120 125 130															
aat tca aca tcg acc tta tcc act tgt tta att aat caa att tta tca															546
Asn Ser Thr Ser Thr Leu Ser Thr Cys Leu Ile Asn Gln Ile Leu Ser															
135 140 145															
ctc aat aga gca tca cct gag ata gtg gga aaa ggt tgt tta aag gaa															594
Leu Asn Arg Ala Ser Pro Glu Ile Val Gly Lys Gly Cys Leu Lys Glu															
150 155 160 165															



tct ggg tca tac atg tgg ata tat gtg ttc atg ggt aat atg ctt cgt Ser Gly Ser Tyr Met Trp Ile Tyr Val Phe Met Gly Asn Met Leu Arg 170 175 180	642
gga ata ggg gag act ccc ata gta cca ytg ggg ctt tct tac att gat Gly Ile Gly Glu Thr Pro Ile Val Pro Xaa Gly Leu Ser Tyr Ile Asp 185 190 195	690
gat ttc gct aaa gaa gga cat tct tct ttg tat tta ggt ata ttg aat Asp Phe Ala Lys Glu Gly His Ser Ser Leu Tyr Leu Gly Ile Leu Asn 200 205 210	738
gca ata gca atg att ggt cca atc att ggc ttt acc ctg gga tct ctg Ala Ile Ala Met Ile Gly Pro Ile Ile Gly Phe Thr Leu Gly Ser Leu 215 220 225	786
ttt tct aaa atg tac gtg gat att gga tat gta gat cta agc act atc Phe Ser Lys Met Tyr Val Asp Ile Gly Tyr Val Asp Leu Ser Thr Ile 230 235 240 245	834
agg ata act cct act gat tct cga tgg gtt gga gct tgg tgg ctt aat Arg Ile Thr Pro Thr Asp Ser Arg Trp Val Gly Ala Trp Trp Leu Asn 250 255 260	882
ttc ctt gtg tct gga cta ttc tcc att att tct tcc ata cca ttc ttt Phe Leu Val Ser Gly Leu Phe Ser Ile Ile Ser Ser Ile Pro Phe Phe 265 270 275	930
ttc ttg ccc caa act cca aat aaa cca caa aaa gaa aga aaa gct tca Phe Leu Pro Gln Thr Pro Asn Lys Pro Gln Lys Glu Arg Lys Ala Ser 280 285 290	978
ctg tct ttg cat gtg ctg gaa aca aat gat gaa aag gat caa aca gct Leu Ser Leu His Val Leu Glu Thr Asn Asp Glu Lys Asp Gln Thr Ala 295 300 305	1026
aat ttg acc aat caa gga aaa aat att acc aaa aat gtg act ggt ttt Asn Leu Thr Asn Gln Gly Lys Asn Ile Thr Lys Asn Val Thr Gly Phe 310 315 320 325	1074
ttc cag tct ttt aaa agc atc ctt act aat ccc ctg tat gtt atg ttt Phe Gln Ser Phe Lys Ser Ile Leu Thr Asn Pro Leu Tyr Val Met Phe 330 335 340	1122
gtg ctt ttg acg ttg tta caa gta agc agc tat att ggt gct ttt act Val Leu Leu Thr Leu Leu Gln Val Ser Ser Tyr Ile Gly Ala Phe Thr 345 350 355	1170
tat gtc ttc aaa tac gta gag caa cag tat ggt cag cct tca tct aag Tyr Val Phe Lys Tyr Val Glu Gln Gln Tyr Gly Gln Pro Ser Ser Lys 360 365 370	1218
gct aac atc tta ttg gga gtc ata acc ata cct att ttt gca agt gga Ala Asn Ile Leu Leu Gly Val Ile Thr Ile Pro Ile Phe Ala Ser Gly 375 380 385	1266
atg ttt tta gga gga tat atc att aaa aaa ttc aaa ctg aac acc gtt Met Phe Leu Gly Gly Tyr Ile Ile Lys Lys Phe Lys Leu Asn Thr Val 390 395 400 405	1314

gga att gcc aaa ttc tca tgt ttt act gct gtg atg tca ttg tcc ttt 1362  
 Gly Ile Ala Lys Phe Ser Cys Phe Thr Ala Val Met Ser Leu Ser Phe  
 410 415 420

tac cta tta tat ttt ttc ata ctc tgt gaa aac aaa tca gtt gcc gga 1410  
 Tyr Leu Leu Tyr Phe Phe Ile Leu Cys Glu Asn Lys Ser Val Ala Gly  
 425 430 435

cta acc atg acc tat gat gga aat aat cca gtg aca tct cat aga gat 1458  
 Leu Thr Met Thr Tyr Asp Gly Asn Asn Pro Val Thr Ser His Arg Asp  
 440 445 450

gta cca ctt tct tat tgc aac tca gac tgc aat tgt gat gaa agt caa 1506  
 Val Pro Leu Ser Tyr Cys Asn Ser Asp Cys Asn Cys Asp Glu Ser Gln  
 455 460 465

tgg gaa cca gtc tgt gga aac aat gga ata act tac atc tca ccc tgt 1554  
 Trp Glu Pro Val Cys Gly Asn Asn Gly Ile Thr Tyr Ile Ser Pro Cys  
 470 475 480 485

cta gca ggt tgc aaa tct tca agt ggc aat aaa aag cct ata gtg ttt 1602  
 Leu Ala Gly Cys Lys Ser Ser Ser Gly Asn Lys Lys Pro Ile Val Phe  
 490 495 500

tac aac tgc agt tgt ttg gaa gta act ggt ctc cag aac aga aat tac 1650  
 Tyr Asn Cys Ser Cys Leu Glu Val Thr Gly Leu Gln Asn Arg Asn Tyr  
 505 510 515

tca gcc cat ttg ggt gaa tgc cca aga gat gat gct tgt aca agg aaa 1698  
 Ser Ala His Leu Gly Glu Cys Pro Arg Asp Asp Ala Cys Thr Arg Lys  
 520 525 530

ttt tac ttt ttt gtt gca ata caa gtc ttg aat tta ttt ttc tct gca 1746  
 Phe Tyr Phe Phe Val Ala Ile Gln Val Leu Asn Leu Phe Phe Ser Ala  
 535 540 545

ctt gga ggc acc tca cat gtc atg ctg att gtt aaa att gtt caa cct 1794  
 Leu Gly Gly Thr Ser His Val Met Leu Ile Val Lys Ile Val Gln Pro  
 550 555 560 565

gaa ttg aaa tca ctt gca ctg ggt ttc cac tca atg gtt ata cga gca 1842  
 Glu Leu Lys Ser Leu Ala Leu Gly Phe His Ser Met Val Ile Arg Ala  
 570 575 580

cta gga gga att cta gct cca ata tat ttt ggg gct ctg att gat aca 1890  
 Leu Gly Gly Ile Leu Ala Pro Ile Tyr Phe Gly Ala Leu Ile Asp Thr  
 585 590 595

acg tgt ata aag tgg tcc acc aac aac tgt ggc aca cgt ggg tca tgt 1938  
 Thr Cys Ile Lys Trp Ser Thr Asn Asn Cys Gly Thr Arg Gly Ser Cys  
 600 605 610

agg aca tat aat tcc aca tca ttt tca agg gtc tac ttg ggc ttg tct 1986  
 Arg Thr Tyr Asn Ser Thr Ser Phe Ser Arg Val Tyr Leu Gly Leu Ser  
 615 620 625

tca atg tta aga gtc tca tca ctt gtt tta tat att ata tta att tat 2034  
 Ser Met Leu Arg Val Ser Ser Leu Val Leu Tyr Ile Ile Leu Ile Tyr  
 630 635 640 645

gcc atg aag aaa aaa tat caa gag aaa gat atc aat gca tca gaa aat 2082  
 Ala Met Lys Lys Lys Tyr Gln Glu Lys Asp Ile Asn Ala Ser Glu Asn  
                   650                  655                  660

gga agt gtc atg gat gaa gca aac tta gaa tcc tta aat aaa aat aaa 2130  
 Gly Ser Val Met Asp Glu Ala Asn Leu Glu Ser Leu Asn Lys Asn Lys  
                   665                  670                  675

cat ttt gtc cct tct gct ggg gca gat agt gaa aca cat tgt taa 2175  
 His Phe Val Pro Ser Ala Gly Ala Asp Ser Glu Thr His Cys \*  
                   680                  685                  690

ggggagaaaa aaagccactt ctgcttctgt gtttccaaac agcattgcat tgattcagta 2235  
 agatgttatt tttagaggagt tcctggctct ttctaaga atttccacat cttttatggt 2295  
 ggaagtataa ataagcctat gaacttataa taaaacaaac tgtaggtaga aaaaaaaaaa 2355  
 aaaaaa 2361

<210> 10  
 <211> 691  
 <212> PRT  
 <213> H. sapiens

<220>  
 <221> VARIANT  
 <222> (130)...(130)  
 <223> Xaa = Asp or Asn

<221> VARIANT  
 <222> (191)...(191)  
 <223> Xaa = Leu

<400> 10  
 Met Asp Gln Asn Gln His Leu Asn Lys Thr Ala Glu Ala Gln Pro Ser  
   1                  5                  10                  15  
 Glu Asn Lys Lys Thr Arg Tyr Cys Asn Gly Leu Lys Met Phe Leu Ala  
                   20                  25                  30  
 Ala Leu Ser Leu Ser Phe Ile Ala Lys Thr Leu Gly Ala Ile Ile Met  
                   35                  40                  45  
 Lys Ser Ser Ile Ile His Ile Glu Arg Arg Phe Glu Ile Ser Ser Ser  
                   50                  55                  60  
 Leu Val Gly Phe Ile Asp Gly Ser Phe Glu Ile Gly Asn Leu Leu Val  
   65                  70                  75                  80  
 Ile Val Phe Val Ser Tyr Phe Gly Ser Lys Leu His Arg Pro Lys Leu  
                   85                  90                  95  
 Ile Gly Ile Gly Cys Phe Ile Met Gly Ile Gly Gly Val Leu Thr Ala  
                   100                  105                  110  
 Leu Pro His Phe Phe Met Gly Tyr Tyr Arg Tyr Ser Lys Glu Thr Asn  
                   115                  120                  125  
 Ile Xaa Ser Ser Glu Asn Ser Thr Ser Thr Leu Ser Thr Cys Leu Ile  
                   130                  135                  140  
 Asn Gln Ile Leu Ser Leu Asn Arg Ala Ser Pro Glu Ile Val Gly Lys  
   145                  150                  155                  160  
 Gly Cys Leu Lys Glu Ser Gly Ser Tyr Met Trp Ile Tyr Val Phe Met  
                   165                  170                  175  
 Gly Asn Met Leu Arg Gly Ile Gly Glu Thr Pro Ile Val Pro Xaa Gly  
                   180                  185                  190  
 Leu Ser Tyr Ile Asp Asp Phe Ala Lys Glu Gly His Ser Ser Leu Tyr  
                   195                  200                  205  
 Leu Gly Ile Leu Asn Ala Ile Ala Met Ile Gly Pro Ile Ile Gly Phe  
                   210                  215                  220  
 Thr Leu Gly Ser Leu Phe Ser Lys Met Tyr Val Asp Ile Gly Tyr Val  
   225                  230                  235                  240

Asp Leu Ser Thr Ile Arg Ile Thr Pro Thr Asp Ser Arg Trp Val Gly  
 245 250 255  
 Ala Trp Trp Leu Asn Phe Leu Val Ser Gly Leu Phe Ser Ile Ile Ser  
 260 265 270  
 Ser Ile Pro Phe Phe Phe Leu Pro Gln Thr Pro Asn Lys Pro Gln Lys  
 275 280 285  
 Glu Arg Lys Ala Ser Leu Ser Leu His Val Leu Glu Thr Asn Asp Glu  
 290 295 300  
 Lys Asp Gln Thr Ala Asn Leu Thr Asn Gln Gly Lys Asn Ile Thr Lys  
 305 310 315 320  
 Asn Val Thr Gly Phe Phe Gln Ser Phe Lys Ser Ile Leu Thr Asn Pro  
 325 330 335  
 Leu Tyr Val Met Phe Val Leu Leu Thr Leu Leu Gln Val Ser Ser Tyr  
 340 345 350  
 Ile Gly Ala Phe Thr Tyr Val Phe Lys Tyr Val Glu Gln Gln Tyr Gly  
 355 360 365  
 Gln Pro Ser Ser Lys Ala Asn Ile Leu Leu Gly Val Ile Thr Ile Pro  
 370 375 380  
 Ile Phe Ala Ser Gly Met Phe Leu Gly Gly Tyr Ile Ile Lys Lys Phe  
 385 390 395 400  
 Lys Leu Asn Thr Val Gly Ile Ala Lys Phe Ser Cys Phe Thr Ala Val  
 405 410 415  
 Met Ser Leu Ser Phe Tyr Leu Leu Tyr Phe Phe Ile Leu Cys Glu Asn  
 420 425 430  
 Lys Ser Val Ala Gly Leu Thr Met Thr Tyr Asp Gly Asn Asn Pro Val  
 435 440 445  
 Thr Ser His Arg Asp Val Pro Leu Ser Tyr Cys Asn Ser Asp Cys Asn  
 450 455 460  
 Cys Asp Glu Ser Gln Trp Glu Pro Val Cys Gly Asn Asn Gly Ile Thr  
 465 470 475 480  
 Tyr Ile Ser Pro Cys Leu Ala Gly Cys Lys Ser Ser Ser Gly Asn Lys  
 485 490 495  
 Lys Pro Ile Val Phe Tyr Asn Cys Ser Cys Leu Glu Val Thr Gly Leu  
 500 505 510  
 Gln Asn Arg Asn Tyr Ser Ala His Leu Gly Glu Cys Pro Arg Asp Asp  
 515 520 525  
 Ala Cys Thr Arg Lys Phe Tyr Phe Phe Val Ala Ile Gln Val Leu Asn  
 530 535 540  
 Leu Phe Phe Ser Ala Leu Gly Gly Thr Ser His Val Met Leu Ile Val  
 545 550 555 560  
 Lys Ile Val Gln Pro Glu Leu Lys Ser Leu Ala Leu Gly Phe His Ser  
 565 570 575  
 Met Val Ile Arg Ala Leu Gly Gly Ile Leu Ala Pro Ile Tyr Phe Gly  
 580 585 590  
 Ala Leu Ile Asp Thr Thr Cys Ile Lys Trp Ser Thr Asn Asn Cys Gly  
 595 600 605  
 Thr Arg Gly Ser Cys Arg Thr Tyr Asn Ser Thr Ser Phe Ser Arg Val  
 610 615 620  
 Tyr Leu Gly Leu Ser Ser Met Leu Arg Val Ser Ser Leu Val Leu Tyr  
 625 630 635 640  
 Ile Ile Leu Ile Tyr Ala Met Lys Lys Lys Tyr Gln Glu Lys Asp Ile  
 645 650 655  
 Asn Ala Ser Glu Asn Gly Ser Val Met Asp Glu Ala Asn Leu Glu Ser  
 660 665 670  
 Leu Asn Lys Asn Lys His Phe Val Pro Ser Ala Gly Ala Asp Ser Glu  
 675 680 685  
 Thr His Cys  
 690

<210> 11  
 <211> 23  
 <212> DNA

<213> H. sapiens  
<400> 11  
ggggctctga ttgatacaac gtg 23  
<210> 12  
<211> 30  
<212> DNA  
<213> H. sapiens  
<400> 12  
actgtggcac acgtgggtca tgtaggacat 30  
<210> 13  
<211> 20  
<212> DNA  
<213> H. sapiens  
<400> 13  
ctgctgcaa ctaacattgc 20  
<210> 14  
<211> 20  
<212> DNA  
<213> H. sapiens  
<400> 14  
cacacactaa ccatgcctct 20  
<210> 15  
<211> 20  
<212> DNA  
<213> H. sapiens  
<400> 15  
tccagtcatt ggctttgcac 20  
<210> 16  
<211> 23  
<212> DNA  
<213> H. sapiens  
<400> 16  
aagaaccaat aaagctgctt act 23  
<210> 16  
<211> 20  
<212> DNA  
<213> H. sapiens  
<400> 16  
gtgtttgcta gccaccttga 20  
<210> 17  
<211> 20  
<212> DNA  
<213> H. sapiens  
<400> 17  
ggcaacactt cctcaaagtg 20  
<210> 18

<211> 20  
<212> DNA  
<213> H. sapiens

<400> 18 20  
gatgctttcc tctgtgcagt

<210> 19  
<211> 20  
<212> DNA  
<213> H. sapiens

<400> 19 20  
ccttcaagcc gaagaaggct

<210> 20  
<211> 20  
<212> DNA  
<213> H. sapiens

<400> 20 20  
aggagttcct ggtcctttca

<210> 21  
<211> 20  
<212> DNA  
<213> H. sapiens

<400> 21 20  
caagctagac ttcaggcctt

<210> 22  
<211> 24  
<212> DNA  
<213> H. sapiens

<400> 22 24  
gaggaattct agtccaata tatt

<210> 23  
<211> 21  
<212> DNA  
<213> H. sapiens

<400> 23 21  
gtcctacatg acccacgtgt g

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>C12N 15/12, 5/10, C12Q 1/68, A01K 67/00, C12N 15/00, C07K 16/28, 14/705</b>		<b>A3</b>	(11) International Publication Number: <b>WO 00/08157</b> (43) International Publication Date: 17 February 2000 (17.02.00)
(21) International Application Number: <b>PCT/US99/17823</b> (22) International Filing Date: 6 August 1999 (06.08.99) (30) Priority Data: 60/095,835 7 August 1998 (07.08.98) <b>US</b> (71) Applicant (for all designated States except US): <b>AXYS PHARMACEUTICALS, INC.</b> [US/US]; 180 Kimball Way, South San Francisco, CA 94080 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): <b>LAUBERT, Boris</b> [US/US]; 4550 Bancroft Street #4, San Diego, CA 92116 (US). <b>CARDOSO, Gizela</b> [US/US]; 721 North Main Street, Brockton, Maine 02401 (US). <b>HU, Ping</b> [US/US]; 5807 Folkstone Road, Bethesda, MD 20817 (US). <b>MILLER, Andrew, P.</b> [US/US]; 3271 Countryside Drive, San Mateo, CA 94403 (US). <b>BUCKLER, Alan, J.</b> [US/US]; 2315 Lagoon View Drive, Cardiff, CA 92007 (US). (74) Agent: <b>SHERWOOD, Pamela, J.; Bozicevic, Field &amp; Francis LLP</b> , Suite 200, 285 Hamilton Avenue, Palo Alto, CA 94301 (US).		(81) Designated States: <b>AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</b> <b>Published</b> <i>With international search report.</i> (88) Date of publication of the international search report: 23 November 2000 (23.11.00)	
(54) Title: <b>HUMAN ANION TRANSPORTER GENES ATNOV</b>			
(57) Abstract <p>Methods for isolating ATnov genes are provided. The ATnov nucleic acid compositions find use in identifying homologous or related proteins and the DNA sequences encoding such proteins; in producing compositions that modulate the expression or function of the protein; and in studying associated physiological pathways. In addition, modulation of the gene activity in vivo is used for prophylactic and therapeutic purposes, such as identification of cell type based on expression, and the like.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		



# INTERNATIONAL SEARCH REPORT

International Application No

PL./US 99/17823

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C12N15/12 C12N5/10 C12Q1/68 A01K67/00 C12N15/00  
C07K16/28 C07K14/705

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K A61K C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	KULLAK-UBLICK ET AL: "Molecular and functional characterization of an organic anion transporting polypeptide cloned from human liver" GASTROENTEROLOGY, vol. 109, 1 October 1995 (1995-10-01), pages 1274-1282, XP002092892 ISSN: 0016-5085 cited in the application the whole document	1
A	WO 97 31111 A (INTROGENE BV ;AMC AMSTERDAM (NL); KANKER INST NL (NL); OUDE ELFERI) 28 August 1997 (1997-08-28) the whole document --- -/--	1

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*Z\* document member of the same patent family

Date of the actual completion of the international search

9 March 2000

Date of mailing of the international search report

11 3. 07. 00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

CHAMBONNET, F

## INTERNATIONAL SEARCH REPORT

International Application No

PC./US 99/17823

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	Embase accession number AB026257.1 Created 08-JUN-1999 , Sequence characterization: Homo sapiens mRNA for organic anion transporter OATP-C, complete XP002132655 the whole document	1-3
T	--- HSIANG B.H., ZHU Y., WANG Z., WU Y., SASSEVILLE V., YANG W.-P.,: "A novel human hepatic organic anion transporting polypeptide (OATP2) Identification of a liver-specific human organic anion transporting polypeptide and identification of rat and human hydroxymethylglutaryl-coa reductase inhibitor transporter." JOURNAL OF BIOLOGICAL CHEMISTRY (MICROFILMS), vol. 274, no. 52, 24 December 1999 (1999-12-24), pages 37161-37168, XP002131865 MD US the whole document	1-3
T	--- ABE T., ET AL.: "Identification of a Novel Gene Family Encoding Human Liver-specific Organic Anion Transporter LST-1" J. BIOL. CHEM., vol. 274, no. 24, 11 June 1999 (1999-06-11), pages 17159-17163, XP000877288 the whole document -----	1-4

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 99/17823

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Claims 2, 3, 15-17 completely, and 1, 4-14 partially.

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 2 3 15-17 and partially 1 4-14

An isolated nucleic acid comprising a nucleotide sequence selected from the group of SEQ NO:3, SEQ NO:5, SEQ NO:7, and SEQ NO:9, or encoding a mammalian ATnov3 protein, having an amino acid sequence selected from the group of SEQ NO:4, SEQ NO:6, SEQ NO:8 and SEQ NO:10; an expression cassette comprising it; a cell comprising said expression cassette; a method for producing a protein using said cell; mammalian ATnov3 polypeptide; a monoclonal antibody binding to said protein; a non human transgenic animal model for ATnov3 gene function; an isolated nucleic acid probe comprising an ATnov3 sequence polymorphism; an array of oligonucleotides comprising probes for detection of ATnov3 locus polymorphisms.

2. Claims: partially 1 4-14

An isolated nucleic acid comprising the nucleotide sequence as set forth in SEQ NO:1 or encoding a mammalian ATnov1 protein; an expression cassette comprising it; a cell comprising said expression cassette; a method for producing a protein using said cell; mammalian ATnov1 polypeptide; a monoclonal antibody binding to said protein; a non human transgenic animal model for ATnov1 gene function;

3. Claims: partially 1 4-14

An isolated nucleic acid comprising the nucleotide sequence as set forth in SEQ NO:2 or encoding a mammalian ATnov2 protein; an expression cassette comprising it; a cell comprising said expression cassette; a method for producing a protein using said cell; mammalian ATnov2 polypeptide; a monoclonal antibody binding to said protein; a non human transgenic animal model for ATnov2 gene function;

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PL /US 99/17823

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9731111 A	28-08-1997	AU 1736697 A	10-09-1997
		EP 0900273 A	10-03-1999
-----			